arXiv:1406.6318v1 [math-ph] 24 Jun 2014

SUSY gauge theory on graded manifolds

G. Sardanashvily, W. Wachowski

Department of Theoretical Physics, Moscow State University, Moscow, Russia

Abstract

Lagrangian classical field theory of even and odd fields is adequately formulated in terms of fibre bundles and graded manifolds. In particular, conventional Yang–Mills gauge theory is theory of connections on smooth principal bundles, but its BRST extension involves odd ghost fields an antifields on graded manifolds. Here, we formulate Yang–Mills theory of Grassmann-graded gauge fields associated to Lie superalgebras on principal graded bundles. A problem lies in a geometric definition of odd gauge fields. Our goal is Yang–Mills theory of graded gauge fields and its BRST extension.

1 Introduction

Conventional Yang–Mills theory of classical gauge fields is adequately formulated as Lagrangian theory of principal connections on smooth principal bundles [20, 25]. Here, we aim to develop Yang–Mills theory of Grassmann-graded even and odd gauge fields associated to Lie superalgebras which characterize various SUSY extensions of Standard Model. A key point is the geometric description of odd gauge fields. A main contradiction is that gauge fields are affine objects, whereas odd fields are linear. A problem also lies in definition of odd fields and their jets.

Even classical fields on an n-dimensional smooth manifold X are adequately represented by sections of some fibre bundle $Y \to X$, and their Lagrangian theory is comprehensively formulated in terms of the variational bicomplex on jet manifolds J^*Y of Y [20, 33]. This formulation is based on the categorial equivalence of projective $C^\infty(X)$-modules of finite ranks and vector bundles over X in accordance with the well-known Serre–Swan theorem, generalized to an arbitrary manifold [20, 28].

At the same time, different geometric models of odd variables either on graded manifolds or supermanifolds are discussed [12, 13, 14, 26, 27, 34]. Both graded manifolds and supermanifolds are phrased in terms of sheaves of graded commutative algebras [4, 20, 31]. However, graded manifolds are characterized by sheaves on smooth manifolds, while supermanifolds are constructed by gluing of sheaves on supervector spaces. Treating odd fields on a smooth manifold X, we follow the above mentioned Serre–Swan theorem extended to graded manifolds (Theorem 2) [34, 35]. It states that, if a graded commutative $C^\infty(X)$-ring is generated by a projective $C^\infty(X)$-module of finite rank, it is isomorphic to a ring of graded functions on a graded manifold whose body is X. In accordance with this theorem, we describe odd fields in terms of graded manifolds [20, 33, 34].

Let us recall that a graded manifold is a locally-ringed space, characterized by a smooth body manifold Z and some structure sheaf $\mathfrak{A}$ of Grassmann algebras on Z [4, 20, 31]. This sheaf is projected as σ (19) onto a sheaf C^∞_Z of smooth real functions on Z. Its sections form a graded commutative $C^\infty(Z)$-ring $\mathcal{A}$ of graded functions on a graded manifold $(Z, \mathfrak{A})$. It is

called the structure ring of $(Z, \mathfrak{A})$. The differential calculus on a graded manifold is defined as the Chevalley–Eilenberg differential calculus on its structure ring (Section 3). However, it is not a particular noncommutative differential calculus because the Leibniz rule (9) of derivations of a graded commutative ring differs from that of a noncommutative ring [33].

It is important for our consideration that, by virtue of the well-known Batchelor theorem (Theorem 1), there exists a vector bundle $E \to Z$ with a typical fibre V such that the structure sheaf $\mathfrak{A}$ of $(Z, \mathfrak{A})$ is isomorphic to a sheaf $\mathfrak{A}_E$ of germs of sections of the exterior bundle $\wedge E^*$ of the dual E^* of E whose typical fibre is the Grassmann algebra $\wedge V^*$ [4, 8]. This Batchelor's isomorphism is not canonical. In field models, it however is fixed from the beginning. Therefore, we restrict our consideration to graded manifolds $(Z, \mathfrak{A}_E)$, called the simple graded manifolds, modelled over vector bundles $E \to Z$.

Let us note that a smooth manifold Z itself can be treated as a trivial graded manifold (Z, C^∞_Z) whose structure ring of graded functions is reduced to a ring $C^\infty(Z)$ of smooth real functions on Z (Remark 1). Accordingly, a fibre bundle $Y \to X$ in a theory of even classical fields can be regarded as a graded bundle of trivial graded manifolds $(Y, C^\infty_Y) \to (X, C^\infty_X)$ (Remark 2). It follows that, in a general setting, one can define a configuration space of theory of even and odd classical fields as being a graded bundle of graded manifolds $(Y, \mathfrak{A}) \to (X, \mathfrak{A}')$, where $Y \to X$ is a smooth bundle (Definition 3).

Considering even and odd fields on a smooth manifold X, we however restrict our analysis to graded bundles

$$(Y, \mathfrak{A}) \to (X, C^\infty_X). \tag{1}$$

over a trivial graded manifold (X, C^∞_X) (Definition 4). We call it the graded bundle over a smooth manifold X. It is denoted by $(X, Y, \mathfrak{A})$. If $Y \to X$ is a vector bundle, this is a particular case of graded vector bundles in [16, 27] whose base is a trivial graded manifold. Accordingly, (X, Y, C^∞_Y) denotes a fibre bundle $Y \to X$.

As was mentioned above, Lagrangian field theory on a fibre bundle $Y \to X$ is formulated in terms of the variational bicomplex on jet manifolds J^*Y of Y. These are fibre bundles over X and, therefore, they can be regarded as trivial graded bundles $(X, J^kY, C^\infty_{J^kY})$. Then let us define their partners in the case of the non-trivial graded manifold (1) as follows (Section 6).

Let $(Y, \mathfrak{A}_F)$ (1) be a simple graded manifold modelled over a vector bundle $F \to Y$ where $Y \to X$ is a fibre bundle. We have a composite bundle $F \to Y \to X$. Let us consider a k-order jet manifold J^kF of $F \to X$. This is a composite bundle $J^kF \to J^kY \to X$ which is a vector bundle $J^kF \to J^kY$ over a k-order jet manifold J^kY of $Y \to X$. Then one can consider a simple graded manifold $(J^kY, \mathfrak{A}_{J^kF})$ which also is a graded bundle $(X, J^kY, \mathfrak{A}_{J^kF})$ over a smooth manifold X. We agree to call it the graded k-order jet manifold of a graded bundle $(X, Y, \mathfrak{A}_F)$ (Definition 7). Obviously, the above mentioned trivial graded jet manifold $(X, J^kY, C^\infty_{J^kY})$ of $(Y, \mathfrak{A}_{F=Y\times\{0\}} = C^\infty_Y)$ is a trivial graded bundle [20, 34].

This definition differs from that of graded jet bundles in [16, 27], but it reproduces the heuristic notion of jets of odd ghosts in field-antifield BRST theory [3, 10].

Furthermore, the inverse sequence of jet manifolds

$$Y \xleftarrow{\pi} J^1Y \longleftarrow \cdots J^{r-1}Y \xleftarrow{\pi^r_{r-1}} J^rY \longleftarrow \cdots, \tag{2}$$

yields the inverse sequence of graded jet manifolds

$$(Y,\mathcal{A}_F) \longleftarrow (J^1Y,\mathfrak{A}_{J^1F}) \longleftarrow \cdots (J^{r-1}Y,\mathfrak{A}_{J^{r-1}F}) \longleftarrow (J^rY,\mathfrak{A}_{J^rF}) \longleftarrow \cdots . \tag{3}$$

One can think on its inverse limit $(J^\infty Y,\mathcal{A}_{J^\infty F})$ as the graded infinite order jet manifold whose body is an infinite order jet manifold $J^\infty Y$ and whose structure sheaf $\mathcal{A}_{J^\infty F}$ is a sheaf of germs of graded functions on graded manifolds $(J^*Y,\mathfrak{A}_{J^*F})$. However $(J^\infty Y,\mathcal{A}_{J^\infty F})$ fails to be a graded manifold in a strict sense because the inverse image $J^\infty Y$ of the sequence (2) is a Fréche manifold, but not the smooth one.

Classical Lagrangian theory of even and odd fields is comprehensively formulated in terms of a Grassmann-graded variational bicomplex on a graded infinite order jet manifold $(J^\infty Y,\mathcal{A}_{J^\infty F})$ (Section 7) [20, 34].

However, quantization of Lagrangian field theory essentially depends on its degeneracy characterized by a set of non-trivial reducible Noether identities [3, 7, 21]. These Noether identities can obey first-stage Noether identities, which in turn are subject to the second-stage ones, and so on. Thus, there is a hierarchy of Noether and higher-stage Noether identities [7, 20]. Bearing in mind gauge theory, we here restrict our consideration to irreducible Lagrangian systems whose Noether identities are independent (Section 8). The inverse second Noether theorem (Theorem 16) associates to them the gauge operator (74) which is a non-trivial gauge symmetry of an original Lagrangian. Its nilpotent extension is the BRST operator (77). If the BRST operator exists, it provides the BRST extension of original Lagrangian field theory by Grassmann-graded ghosts and antifields [19, 20]. This BRST extension is a first step towards BV quantization of classical field theory in terms of functional integrals [3, 21].

As was mentioned above, classical gauge theory is formulated as Lagrangian theory of principal connections on smooth principal bundles (Section 9). Similarly, we develop SUSY gauge theory in the framework of Grassmann-graded Lagrangian formalism on principal graded bundles (Section 10). A key point is that we consider simple principal graded bundles (Definition 26) subject to an action of an even Lie group, but not the whole graded Lie one. In this case, odd gauge potentials in comparison with the even ones are linear, but not affine objects (see Remark 10). At the same time, they admit the affine gauge transformations (141) parameterised by ghosts. Our goal is Yang–Mills theory of graded gauge fields and its BRST extension (Section 11).

2 Grassmann-graded algebraic calculus

Throughout this work, by the Grassmann gradation is meant $\mathbb{Z}_2$-gradation, and a Grassmann graded structure simply is called the graded structure if there is no danger of confusion. Hereafter, the symbol [.] stands for the Grassmann parity. Let us summarize the relevant notions of the Grassmann-graded algebraic calculus [4, 20, 31].

Let $\mathcal{K}$ be a commutative ring. A $\mathcal{K}$-module Q is called graded if it is endowed with a grading automorphism γ, $\gamma^2=\mathrm{Id}$. A graded module falls into a direct sum of modules $Q=Q_0\oplus Q_1$ such that

$$\gamma(q)=(-1)^{[q]}q, \qquad q\in Q_{[q]}.$$

One calls Q_0 and Q_1 the even and odd parts of Q, respectively. A graded $\mathcal{K}$-module is said to be free if it has a basis composed by graded-homogeneous elements.

In particular, by a real graded vector space $B = B_0 \oplus B_1$ is meant a graded $\mathbb{R}$-module. A real graded vector space is said to be (n, m)-dimensional if $B_0 = \mathbb{R}^n$ and $B_1 = \mathbb{R}^m$.

A $\mathcal{K}$-algebra $\mathcal{A}$ is called graded if it is a graded $\mathcal{K}$-module such that

$$[aa'] = ([a] + [a'])\mathrm{mod}\, 2,$$

where a and a' are graded-homogeneous elements of $\mathcal{A}$. Its even part $\mathcal{A}_0$ is a subalgebra of $\mathcal{A}$, and the odd one $\mathcal{A}_1$ is an $\mathcal{A}_0$-module. If $\mathcal{A}$ is a graded ring, then $[\mathbf{1}] = 0$.

Given a graded algebra $\mathcal{A}$, a left graded $\mathcal{A}$-module Q is defined as a left $\mathcal{A}$-module where

$$[aq] = ([a] + [q])\mathrm{mod}\, 2.$$

Similarly, right graded $\mathcal{A}$-modules are treated.

A graded algebra $\mathcal{A}$ is called graded commutative if

$$aa' = (-1)^{[a][a']}a'a.$$

In this case, a graded $\mathcal{A}$-module Q is provided with a graded $\mathcal{A}$-bimodule structure by letting

$$qa = (-1)^{[a][q]}aq, \qquad a \in \mathcal{A}, \qquad q \in Q.$$

Given a graded commutative ring $\mathcal{A}$, the following are standard constructions of new graded modules from old ones.

• The direct sum of graded modules and a graded factor module are defined just as those of modules over a commutative ring.

• The tensor product $P \otimes Q$ of graded $\mathcal{A}$-modules P and Q is their tensor product as $\mathcal{A}$-modules such that

$$\begin{aligned} &[p \otimes q] = [p] + [q], \qquad p \in P, \qquad q \in Q, \\ &ap \otimes q = (-1)^{[p][a]}pa \otimes q = (-1)^{[p][a]}p \otimes aq, \qquad a \in \mathcal{A}. \end{aligned}$$

In particular, the tensor algebra $\otimes P$ of a graded $\mathcal{A}$-module P is defined just as that of a module over a commutative ring. Its quotient $\wedge P$ with respect to the ideal generated by elements

$$p \otimes p' + (-1)^{[p][p']}p' \otimes p, \qquad p, p' \in P,$$

is the bigraded exterior algebra of a graded module P with respect to the graded exterior product

$$p \wedge p' = -(-1)^{[p][p']}p' \wedge p.$$

• A morphism $\Phi : P \to Q$ of graded $\mathcal{A}$-modules seen as additive groups is said to be an even graded morphism (resp. an odd graded morphism) if Φ preserves (resp. changes) the Grassmann parity of all graded-homogeneous elements of P and if the relations

$$\Phi(ap) = (-1)^{[\Phi][a]}a\Phi(p), \qquad p \in P, \qquad a \in \mathcal{A},$$

hold. A morphism $\Phi : P \to Q$ of graded $\mathcal{A}$-modules as additive groups is called a graded $\mathcal{A}$-module morphism if it is represented by a sum of even and odd graded morphisms. A set

$\mathrm{Hom}_{\mathcal{A}}(P,Q)$ of graded morphisms of a graded $\mathcal{A}$-module P to a graded $\mathcal{A}$-module Q is naturally a graded $\mathcal{A}$-module. A graded $\mathcal{A}$-module $P^* = \mathrm{Hom}_{\mathcal{A}}(P,\mathcal{A})$ is called the dual of a graded $\mathcal{A}$-module P.

Let V be a real vector space, and let $\Lambda = \wedge V$ be its exterior algebra endowed with the Grassmann gradation

$$\Lambda = \Lambda_0 \oplus \Lambda_1, \qquad \Lambda_0 = \mathbb{R} \bigoplus_{k=1} \overset{2k}{\wedge} V, \qquad \Lambda_1 = \bigoplus_{k=1} \overset{2k-1}{\wedge} V. \tag{4}$$

It is a real graded commutative ring, called the Grassmann algebra. A Grassmann algebra, seen as an additive group, admits the decomposition

$$\Lambda = \mathbb{R} \oplus R = \mathbb{R} \oplus R_0 \oplus R_1 = \mathbb{R} \oplus (\Lambda_1)^2 \oplus \Lambda_1, \tag{5}$$

where R is the ideal of nilpotents of Λ. The corresponding epimorphism $\sigma : \Lambda \to \mathbb{R}$ is called the body map.

Note that there is a different definition of a Grassmann algebra which is equivalent to the above mentioned one only in the case of infinite-dimensional vector spaces V [22]. Hereafter, we restrict our consideration to Grassmann algebras of finite rank when $V = \mathbb{R}^N$. Given a basis $\{c^i\}$ for V, elements of the Grassmann algebra Λ (4) take a form

$$a = \sum_{k=0,1,\dots} \sum_{(i_1\cdots i_k)} a_{i_1\cdots i_k} c^{i_1}\cdots c^{i_k}, \tag{6}$$

where the second sum runs through all the tuples $(i_1 \cdots i_k)$ such that no two of them are permutations of each other.

A graded algebra $\mathfrak{g}$ is called a Lie superalgebra if its product $[.,.]$, called the Lie superbracket, obeys the relations

$$[\varepsilon,\varepsilon'] = -(-1)^{[\varepsilon][\varepsilon']}[\varepsilon',\varepsilon],$$
$$(-1)^{[\varepsilon][\varepsilon'']}[\varepsilon,[\varepsilon',\varepsilon'']] + (-1)^{[\varepsilon'][\varepsilon]}[\varepsilon',[\varepsilon'',\varepsilon]] + (-1)^{[\varepsilon''][\varepsilon']}[\varepsilon'',[\varepsilon,\varepsilon']] = 0.$$

Being decomposed in even and odd parts $\mathfrak{g} = \mathfrak{g}_0 \oplus \mathfrak{g}_1$, a Lie superalgebra $\mathfrak{g}$ obeys the relations

$$[\mathfrak{g}_0,\mathfrak{g}_0] \subset \mathfrak{g}_0, \qquad [\mathfrak{g}_0,\mathfrak{g}_1] \subset \mathfrak{g}_1, \qquad [\mathfrak{g}_1,\mathfrak{g}_1] \subset \mathfrak{g}_1.$$

In particular, an even part $\mathfrak{g}_0$ of a Lie superalgebra $\mathfrak{g}$ is a Lie algebra.

3 Grassmann-graded differential calculus

Linear differential operators on graded modules over a graded commutative ring are defined similarly to those in commutative geometry [20, 31, 32].

Let $\mathcal{K}$ be a commutative ring and $\mathcal{A}$ a graded commutative $\mathcal{K}$-ring. Let P and Q be graded $\mathcal{A}$-modules. A $\mathcal{K}$-module $\mathrm{Hom}_{\mathcal{K}}(P,Q)$ of graded $\mathcal{K}$-module homomorphisms $\Phi : P \to Q$ can be endowed with the two graded $\mathcal{A}$-module structures

$$(a\Phi)(p) = a\Phi(p), \qquad (\Phi \bullet a)(p) = \Phi(ap), \qquad a \in \mathcal{A}, \quad p \in P, \tag{7}$$

called $\mathcal{A}$- and $\mathcal{A}^\bullet$-module structures, respectively. Let us put

$$\delta_a\Phi = a\Phi - (-1)^{[a][\Phi]}\Phi \bullet a, \qquad a \in \mathcal{A}. \tag{8}$$

An element $\Delta \in \mathrm{Hom}_{\mathcal{K}}(P, Q)$ is said to be a Q-valued graded differential operator of order s on P if $\delta_{a_0} \circ \cdots \circ \delta_{a_s}\Delta = 0$ for any tuple of $s+1$ elements $a_0, \ldots, a_s$ of $\mathcal{A}$. A set $\mathrm{Diff}_s(P, Q)$ of these operators inherits the graded module structures (7).

In particular, zero order graded differential operators coincide with graded $\mathcal{A}$-module morphisms $P \to Q$. A first order graded differential operator Δ satisfies a relation

$$\delta_a \circ \delta_b\, \Delta(p) = ab\Delta(p) - (-1)^{([b]+[\Delta])[a]}b\Delta(ap) - (-1)^{[b][\Delta]}a\Delta(bp) + (-1)^{[b][\Delta]+([\Delta]+[b])[a]} = 0, \qquad a, b \in \mathcal{A}, \quad p \in P.$$

For instance, let $P = \mathcal{A}$. Any zero order Q-valued graded differential operator Δ on $\mathcal{A}$ is defined by its value $\Delta(\mathbf{1})$. Then there is a graded $\mathcal{A}$-module isomorphism

$$\mathrm{Diff}_0(\mathcal{A}, Q) = Q, \qquad Q \ni q \to \Delta_q \in \mathrm{Diff}_0(\mathcal{A}, Q),$$

where Δ_q is given by the equality $\Delta_q(\mathbf{1}) = q$. A first order Q-valued graded differential operator Δ on $\mathcal{A}$ fulfils a condition

$$\Delta(ab) = \Delta(a)b + (-1)^{[a][\Delta]}a\Delta(b) - (-1)^{([b]+[a])[\Delta]}ab\Delta(\mathbf{1}), \qquad a, b \in \mathcal{A}.$$

It is called the Q-valued graded derivation of $\mathcal{A}$ if $\Delta(\mathbf{1}) = 0$, i.e., a graded Leibniz rule

$$\Delta(ab) = \Delta(a)b + (-1)^{[a][\Delta]}a\Delta(b), \quad a, b \in \mathcal{A}, \tag{9}$$

holds. One then observes that any first order graded differential operator on $\mathcal{A}$ falls into a sum

$$\Delta(a) = \Delta(\mathbf{1})a + [\Delta(a) - \Delta(\mathbf{1})a]$$

of a zero order graded differential operator $\Delta(\mathbf{1})a$ and a graded derivation $\Delta(a) - \Delta(\mathbf{1})a$. If ∂ is a graded derivation of $\mathcal{A}$, then $a\partial$ is so for any $a \in \mathcal{A}$. Hence, graded derivations of $\mathcal{A}$ constitute a graded $\mathcal{A}$-module $\mathfrak{d}(\mathcal{A}, Q)$, called the graded derivation module. If $Q = \mathcal{A}$, a graded derivation module $\mathfrak{d}\mathcal{A}$ also is a Lie superalgebra over a commutative ring $\mathcal{K}$ with respect to a superbracket

$$[u, u'] = u \circ u' - (-1)^{[u][u']}u' \circ u, \qquad u, u' \in \mathcal{A}. \tag{10}$$

Since $\mathfrak{d}\mathcal{A}$ is a Lie $\mathcal{K}$-superalgebra, let us consider the Chevalley–Eilenberg complex $C^*[\mathfrak{d}\mathcal{A}; \mathcal{A}]$ where a graded commutative ring $\mathcal{A}$ is a regarded as a $\mathfrak{d}\mathcal{A}$-module [15, 20, 32]. It is the complex

$$0 \to \mathcal{K} \to \mathcal{A} \xrightarrow{d} C^1[\mathfrak{d}\mathcal{A}; \mathcal{A}] \xrightarrow{d} \cdots C^k[\mathfrak{d}\mathcal{A}; \mathcal{A}] \xrightarrow{d} \cdots \tag{11}$$

where

$$C^k[\mathfrak{d}\mathcal{A}; \mathcal{A}] = \mathrm{Hom}_{\mathcal{K}}(\overset{k}{\wedge}\mathfrak{d}\mathcal{A}, \mathcal{A})$$

are $\mathfrak{d}\mathcal{A}$-modules of $\mathcal{K}$-linear graded morphisms of the graded exterior products $\overset{k}{\wedge}\mathfrak{d}\mathcal{A}$ of a graded $\mathcal{K}$-module $\mathfrak{d}\mathcal{A}$ to $\mathcal{A}$. Let us bring homogeneous elements of $\overset{k}{\wedge}\mathfrak{d}\mathcal{A}$ into the form

$$\varepsilon_1 \wedge \cdots \varepsilon_r \wedge \epsilon_{r+1} \wedge \cdots \wedge \epsilon_k, \qquad \varepsilon_i \in \mathfrak{d}\mathcal{A}_0, \quad \epsilon_j \in \mathfrak{d}\mathcal{A}_1.$$

Then the Chevalley–Eilenberg coboundary operator d of the complex (11) is given by the expression

$$\begin{aligned}
&dc(\varepsilon_1\wedge\cdots\wedge\varepsilon_r\wedge\epsilon_1\wedge\cdots\wedge\epsilon_s)= \qquad (12)\\
&\sum_{i=1}^{r}(-1)^{i-1}\varepsilon_i c(\varepsilon_1\wedge\cdots\widehat{\varepsilon_i}\cdots\wedge\varepsilon_r\wedge\epsilon_1\wedge\cdots\epsilon_s)+\\
&\sum_{j=1}^{s}(-1)^{r}\varepsilon_i c(\varepsilon_1\wedge\cdots\wedge\varepsilon_r\wedge\epsilon_1\wedge\cdots\widehat{\epsilon_j}\cdots\wedge\epsilon_s)+\\
&\sum_{1\leq i<j\leq r}(-1)^{i+j}c([\varepsilon_i,\varepsilon_j]\wedge\varepsilon_1\wedge\cdots\widehat{\varepsilon_i}\cdots\widehat{\varepsilon_j}\cdots\wedge\varepsilon_r\wedge\epsilon_1\wedge\cdots\wedge\epsilon_s)+\\
&\sum_{1\leq i<j\leq s}c([\epsilon_i,\epsilon_j]\wedge\varepsilon_1\wedge\cdots\wedge\varepsilon_r\wedge\epsilon_1\wedge\cdots\widehat{\epsilon_i}\cdots\widehat{\epsilon_j}\cdots\wedge\epsilon_s)+\\
&\sum_{1\leq i<r,1\leq j\leq s}(-1)^{i+r+1}c([\varepsilon_i,\epsilon_j]\wedge\varepsilon_1\wedge\cdots\widehat{\varepsilon_i}\cdots\wedge\varepsilon_r\wedge\epsilon_1\wedge\cdots\widehat{\epsilon_j}\cdots\wedge\epsilon_s),
\end{aligned}$$

where the caret $\widehat{\ }$ denotes omission.

It is easily justified that the complex (11) contains a subcomplex $\mathcal{O}^*[\mathfrak{d}\mathcal{A}]$ of $\mathcal{A}$-linear graded morphisms. The $\mathbb{N}$-graded module $\mathcal{O}^*[\mathfrak{d}\mathcal{A}]$ is provided with the structure of a bigraded $\mathcal{A}$-algebra with respect to the graded exterior product

$$\begin{aligned}
&\phi\wedge\phi'(u_1,...,u_{r+s})=\sum_{i_1<\cdots<i_r;j_1<\cdots<j_s}\mathrm{Sgn}_{1\cdots r+s}^{i_1\cdots i_r j_1\cdots j_s}\phi(u_{i_1},\ldots,u_{i_r})\phi'(u_{j_1},\ldots,u_{j_s}), \qquad (13)\\
&\phi\in\mathcal{O}^r[\mathfrak{d}\mathcal{A}], \qquad \phi'\in\mathcal{O}^s[\mathfrak{d}\mathcal{A}], \qquad u_k\in\mathfrak{d}\mathcal{A},
\end{aligned}$$

where $u_1,\ldots,u_{r+s}$ are graded-homogeneous elements of $\mathfrak{d}\mathcal{A}$ and

$$u_1\wedge\cdots\wedge u_{r+s}=\mathrm{Sgn}_{1\cdots r+s}^{i_1\cdots i_r j_1\cdots j_s}u_{i_1}\wedge\cdots\wedge u_{i_r}\wedge u_{j_1}\wedge\cdots\wedge u_{j_s}.$$

The graded Chevalley–Eilenberg coboundary operator d (12) and the graded exterior product $\wedge$ (13) bring $\mathcal{O}^*[\mathfrak{d}\mathcal{A}]$ into a differential bigraded algebra whose elements obey relations

$$\phi\wedge\phi'=(-1)^{|\phi||\phi'|+[\phi][\phi']}\phi'\wedge\phi, \qquad d(\phi\wedge\phi')=d\phi\wedge\phi'+(-1)^{|\phi|}\phi\wedge d\phi'. \qquad (14)$$

It is called the graded Chevalley–Eilenberg differential calculus over a graded commutative $\mathcal{K}$-ring $\mathcal{A}$. In particular, we have

$$\mathcal{O}^1[\mathfrak{d}\mathcal{A}]=\mathrm{Hom}_{\mathcal{A}}(\mathfrak{d}\mathcal{A},\mathcal{A})=\mathfrak{d}\mathcal{A}^*. \qquad (15)$$

One can extend this duality relation to the graded interior product of $u\in\mathfrak{d}\mathcal{A}$ with any element $\phi\in\mathcal{O}^*[\mathfrak{d}\mathcal{A}]$ by the rules

$$\begin{aligned}
&u\rfloor(bda)=(-1)^{[u][b]}bu(a), \qquad a,b\in\mathcal{A},\\
&u\rfloor(\phi\wedge\phi')=(u\rfloor\phi)\wedge\phi'+(-1)^{|\phi|+[\phi][u]}\phi\wedge(u\rfloor\phi'). \qquad (16)
\end{aligned}$$

As a consequence, any graded derivation $u\in\mathfrak{d}\mathcal{A}$ of $\mathcal{A}$ yields a derivation

$$\begin{aligned}
&\mathbf{L}_u\phi=u\rfloor d\phi+d(u\rfloor\phi), \qquad \phi\in\mathcal{O}^*, \qquad u\in\mathfrak{d}\mathcal{A}, \qquad (17)\\
&\mathbf{L}_u(\phi\wedge\phi')=\mathbf{L}_u(\phi)\wedge\phi'+(-1)^{[u][\phi]}\phi\wedge\mathbf{L}_u(\phi'),
\end{aligned}$$

called the graded Lie derivative of the differential bigraded algebra $\mathcal{O}^*[\mathfrak{d}\mathcal{A}]$.

The minimal graded Chevalley–Eilenberg differential calculus $\mathcal{O}^*\mathcal{A} \subset \mathcal{O}^*[\mathfrak{d}\mathcal{A}]$ over a graded commutative ring $\mathcal{A}$ consists of the monomials $a_0 da_1 \wedge \cdots \wedge da_k$, $a_i \in \mathcal{A}$. The corresponding complex

$$0 \to \mathcal{K} \longrightarrow \mathcal{A} \stackrel{d}{\longrightarrow} \mathcal{O}^1\mathcal{A} \stackrel{d}{\longrightarrow} \cdots \mathcal{O}^k\mathcal{A} \stackrel{d}{\longrightarrow} \cdots \tag{18}$$

is called the bigraded de Rham complex of a graded commutative $\mathcal{K}$-ring $\mathcal{A}$.

4 Graded manifolds

A graded manifold of dimension (n, m) is defined as a local-ringed space $(Z, \mathfrak{A})$ where Z is an n-dimensional smooth manifold Z and $\mathfrak{A} = \mathfrak{A}_0 \oplus \mathfrak{A}_1$ is a sheaf of Grassmann algebras Λ of rank m such that [4, 20, 31]:

• there is the exact sequence of sheaves

$$0 \to \mathcal{R} \to \mathfrak{A} \stackrel{\sigma}{\to} C^\infty_Z \to 0, \qquad \mathcal{R} = \mathfrak{A}_1 + (\mathfrak{A}_1)^2, \tag{19}$$

where σ is a body epimorphism onto a sheaf C^∞_Z of smooth real functions on Z;

• $\mathcal{R}/\mathcal{R}^2$ is a locally free sheaf of C^∞_Z-modules of finite rank (with respect to pointwise operations), and the sheaf $\mathfrak{A}$ is locally isomorphic to the exterior product $\wedge_{C^\infty_Z}(\mathcal{R}/\mathcal{R}^2)$.

The sheaf $\mathfrak{A}$ is called a structure sheaf of a graded manifold $(Z, \mathfrak{A})$, and a manifold Z is said to be the body of $(Z, \mathfrak{A})$. Sections of the sheaf $\mathfrak{A}$ are called graded functions on a graded manifold $(Z, \mathfrak{A})$. They make up a graded commutative $C^\infty(Z)$-ring $\mathfrak{A}(Z)$ called the structure ring of $(Z, \mathfrak{A})$.

A graded manifold $(Z, \mathfrak{A})$ possesses the following local structure. Given a point $z \in Z$, there exists its open neighborhood U, called a splitting domain, such that

$$\mathfrak{A}(U) = C^\infty(U) \otimes \wedge\mathbb{R}^m.$$

This means that the restriction $\mathfrak{A}|_U$ of the structure sheaf $\mathfrak{A}$ to U is isomorphic to the sheaf $C^\infty_U \otimes \wedge\mathbb{R}^m$ of sections of some exterior bundle $U \times \wedge\mathbb{R}^m \to U$. The well-known Batchelor theorem [4, 8] states that such a structure of a graded manifold is global as follows.

Theorem 1. Let $(Z, \mathfrak{A})$ be a graded manifold. There exists a vector bundle $E \to Z$ with an m-dimensional typical fibre V such that the structure sheaf $\mathfrak{A}$ of $(Z, \mathfrak{A})$ is isomorphic to the structure sheaf $\mathfrak{A}_E$ of germs of sections of the exterior bundle $\wedge E^* \to Z$, whose typical fibre is a Grassmann algebra $\wedge V^*$.

Combining Batchelor Theorem 1 and the classical Serre–Swan theorem leads to the following Serre–Swan theorem for graded manifolds [6, 35].

Theorem 2. Let Z be a smooth manifold. A graded commutative $C^\infty(Z)$-algebra $\mathcal{A}$ is isomorphic to the structure ring of a graded manifold with a body Z iff it is the exterior algebra of some projective $C^\infty(Z)$-module of finite rank.

It should be emphasized that Batchelor's isomorphism in Theorem 1 fails to be canonical. We agree to call $(Z, \mathfrak{A}_E)$ in Theorem 1 the simple graded manifold modelled over a vector bundle $E \to Z$, named its characteristic vector bundle. Accordingly, a structure ring $\mathfrak{A}_E(Z)$ of a simple graded manifold $(Z, \mathfrak{A}_E)$ is a structure module

$$\mathcal{A}_E = \mathfrak{A}_E(Z) = \wedge E^*(Z) \tag{20}$$

of sections of the exterior bundle $\wedge E^*$.

Remark 1. Hereafter, it is convenient to treat a local-ringed space

$$(Z, \mathfrak{A}_0 = C^\infty_Z) \tag{21}$$

as a trivial graded manifold. It is a simple graded manifold whose characteristic bundle is $E = Z \times \{0\}$. Its structure module is a ring $C^\infty(Z)$ of smooth real functions on Z.

Given a simple graded manifold $(Z, \mathfrak{A}_E)$, every trivialization chart $(U; z^A, q^a)$ of a vector bundle $E \to Z$ yields a splitting domain $(U; z^A, c^a)$ of $(Z, \mathfrak{A}_E)$ where $\{c^a\}$ is the corresponding local fibre basis for $E^* \to X$, i.e., c^a are locally constant sections of $E^* \to X$ such that $q_b \circ c^a = \delta^a_b$. Graded functions on such a chart are Λ-valued functions

$$f = \sum_{k=0}^{m} \frac{1}{k!} f_{a_1 \dots a_k}(z) c^{a_1} \cdots c^{a_k}, \tag{22}$$

where $f_{a_1 \cdots a_k}(z)$ are smooth functions on U. One calls $\{z^A, c^a\}$ the local basis for a graded manifold $(Z, \mathfrak{A}_E)$. Transition functions $q'^a = \rho^a_b(z^A) q^b$ of bundle coordinates on $E \to Z$ induce the corresponding transformation

$$c'^a = \rho^a_b(z^A) c^b \tag{23}$$

of the associated local basis for a graded manifold $(Z, \mathfrak{A}_E)$ and the according coordinate transformation law of graded functions (22).

Note that, in general, automorphisms of a graded manifold take a form $c'^a = \rho^a(z^A, c^b)$, where $\rho^a(z^A, c^b)$ are local graded functions. Considering a simple graded manifold $(Z, \mathfrak{A}_E)$, one restricts the class of graded manifold transformations to the linear ones (23), compatible with given Batchelor's isomorphism.

Let us consider the graded derivation module $\mathfrak{d}\mathfrak{A}(Z)$ of a real graded commutative ring $\mathfrak{A}(Z)$. It is a real Lie superalgebra relative to the superbracket (10). Its elements are called the graded vector fields on a graded manifold $(Z, \mathfrak{A})$. A key point is that graded vector fields $u \in \mathfrak{d}\mathcal{A}_E$ on a simple graded manifold $(Z, \mathfrak{A}_E)$ are represented by sections of some vector bundle as follows [20].

Due to the canonical splitting $VE = E \times E$, the vertical tangent bundle VE of $E \to Z$ can be provided with the fibre bases $\{\partial/\partial c^a\}$, which are the duals of the bases $\{c^a\}$. Then graded vector fields on a splitting domain $(U; z^A, c^a)$ of $(Z, \mathfrak{A}_E)$ read

$$u = u^A \partial_A + u^a \frac{\partial}{\partial c^a}, \tag{24}$$

where u^A, u^a are local graded functions on U. In particular,

$$\frac{\partial}{\partial c^a} \circ \frac{\partial}{\partial c^b} = -\frac{\partial}{\partial c^b} \circ \frac{\partial}{\partial c^a}, \qquad \partial_A \circ \frac{\partial}{\partial c^a} = \frac{\partial}{\partial c^a} \circ \partial_A.$$

The graded derivations (24) act on graded functions $f \in \mathfrak{A}_E(U)$ (22) by the rule

$$u(f_{a \dots b} c^a \cdots c^b) = u^A \partial_A(f_{a \dots b}) c^a \cdots c^b + u^k f_{a \dots b} \frac{\partial}{\partial c^k} \rfloor (c^a \cdots c^b). \tag{25}$$

This rule implies the corresponding coordinate transformation law

$$u'^A = u^A, \qquad u'^a = \rho^a_j u^j + u^A \partial_A(\rho^a_j) c^j$$

of graded vector fields. It follows that graded vector fields (24) are sections of the following vector bundle $\mathcal{V}_E \to Z$. This vector bundle is locally isomorphic to the vector bundle

$$\mathcal{V}_E|_U \approx \wedge E^* \underset{Z}{\otimes} (E \underset{Z}{\oplus} TZ)|_U,$$

and it is characterized by an atlas of bundle coordinates $(z^A, z^A_{a_1\ldots a_k}, v^i_{b_1\ldots b_k})$, $k = 0, \ldots, m$, possessing transition functions

$$z'^A_{i_1\ldots i_k} = \rho^{-1}{}^{a_1}_{i_1} \cdots \rho^{-1}{}^{a_k}_{i_k} z^A_{a_1\ldots a_k},$$
$$v'^i_{j_1\ldots j_k} = \rho^{-1}{}^{b_1}_{j_1} \cdots \rho^{-1}{}^{b_k}_{j_k} \left[\rho^i_j v^j_{b_1\ldots b_k} + \frac{k!}{(k-1)!} z^A_{b_1\ldots b_{k-1}} \partial_A \rho^i_{b_k}\right].$$

Given a structure ring $\mathcal{A}_E$ of graded functions on a simple graded manifold $(Z, \mathfrak{A}_E)$ and the real Lie superalgebra $\mathfrak{d}\mathcal{A}_E$ of its graded derivations, let us consider the graded Chevalley–Eilenberg differential calculus

$$\mathcal{S}^*[E; Z] = \mathcal{O}^*[\mathfrak{d}\mathcal{A}_E] \tag{26}$$

over $\mathcal{A}_E$. Since the graded derivation module $\mathfrak{d}\mathcal{A}_E$ is isomorphic to the structure module of sections of a vector bundle $\mathcal{V}_E \to Z$, elements of $\mathcal{S}^*[E; Z]$ are represented by sections of the exterior bundle $\wedge\overline{\mathcal{V}}_E$ of the $\mathcal{A}_E$-dual $\overline{\mathcal{V}}_E \to Z$ of $\mathcal{V}_E$. With respect to the dual fibre bases $\{dz^A\}$ for T^*Z and $\{dc^b\}$ for E^*, sections of $\overline{\mathcal{V}}_E$ take a coordinate form

$$\phi = \phi_A dz^A + \phi_a dc^a,$$

together with transition functions

$$\phi'_a = \rho^{-1}{}^b_a \phi_b, \qquad \phi'_A = \phi_A + \rho^{-1}{}^b_a \partial_A(\rho^a_j)\phi_b c^j.$$

The duality isomorphism $\mathcal{S}^1[E; Z] = \mathfrak{d}\mathcal{A}^*_E$ (15) is given by the graded interior product

$$u\rfloor\phi = u^A\phi_A + (-1)^{[\phi_a]} u^a \phi_a.$$

Elements of $\mathcal{S}^*[E; Z]$ are called graded exterior forms on a graded manifold $(Z, \mathfrak{A}_E)$.

Seen as an $\mathcal{A}_E$-algebra, the differential bigraded algebra $\mathcal{S}^*[E; Z]$ (26) on a splitting domain $(U; z^A, c^a)$ is locally generated by graded one-forms dz^A, dc^i such that

$$dz^A \wedge dc^i = -dc^i \wedge dz^A, \qquad dc^i \wedge dc^j = dc^j \wedge dc^i.$$

Accordingly, the graded Chevalley–Eilenberg coboundary operator d (12), called the graded exterior differential, reads

$$d\phi = dz^A \wedge \partial_A\phi + dc^a \wedge \frac{\partial}{\partial c^a}\phi,$$

where derivatives ∂_λ, $\partial/\partial c^a$ act on coefficients of graded exterior forms by the formula (25), and they are graded commutative with graded forms dz^A and dc^a. The formulas (14) – (17) hold.

One can show that the differential bigraded algebra $\mathcal{S}^*[E; Z]$ (26) is a minimal differential calculus over $\mathcal{A}_E$, i.e., it is generated by elements df, $f \in \mathcal{A}_E$ [20, 34].

The bigraded de Rham complex (18) of the minimal graded Chevalley–Eilenberg differential calculus $\mathcal{S}^*[E; Z]$ reads

$$0 \to \mathbb{R} \to \mathcal{A}_E \xrightarrow{d} \mathcal{S}^1[E; Z] \xrightarrow{d} \cdots \mathcal{S}^k[E; Z] \xrightarrow{d} \cdots. \tag{27}$$

Its cohomology $H^*(\mathcal{A}_E)$ is called the de Rham cohomology of a simple graded manifold $(Z, \mathfrak{A}_E)$.

In particular, given the differential graded algebra $\mathcal{O}^*(Z)$ of exterior forms on Z, there exists a canonical monomorphism

$$\mathcal{O}^*(Z) \to \mathcal{S}^*[E; Z] \tag{28}$$

and a body epimorphism $\mathcal{S}^*[E; Z] \to \mathcal{O}^*(Z)$ which are cochain morphisms of the de Rham complex (27) and that of $\mathcal{O}^*(Z)$. The de Rham cohomology of a simple graded manifold $(Z, \mathfrak{A}_E)$ equals the de Rham cohomology of its body Z [20].

5 Graded bundles

A morphism of graded manifolds $(Z, \mathfrak{A}) \to (Z', \mathfrak{A}')$ is defined as that of local-ringed spaces

$$\phi : Z \to Z', \qquad \widehat{\Phi} : \mathfrak{A}' \to \phi_* \mathfrak{A}, \tag{29}$$

where ϕ is a manifold morphism and $\widehat{\Phi}$ is a sheaf morphism of $\mathfrak{A}'$ to the direct image $\phi_*\mathfrak{A}$ of $\mathfrak{A}$ onto Z' [17, 38]. The morphism (29) of graded manifolds is said to be:

- a monomorphism if ϕ is an injection and $\widehat{\Phi}$ is an epimorphism;
- an epimorphism if ϕ is a surjection and $\widehat{\Phi}$ is a monomorphism.

Definition 3. An epimorphism of graded manifolds $(Z, \mathfrak{A}) \to (Z', \mathfrak{A}')$ where $Z \to Z'$ is a fibre bundle is called the graded bundle.

Let $E \to Z$ and $E' \to Z$ be vector bundles and $\Phi : E \to E'$ their bundle morphism over a morphism $\phi : Z \to Z'$. Then every section s^* of the dual bundle $E'^* \to Z'$ defines the pull-back section $\Phi^* s^*$ of the dual bundle $E^* \to Z$ by the law

$$v_z \rfloor \Phi^* s^*(z) = \Phi(v_z) \rfloor s^*(\varphi(z)), \qquad v_z \in E_z.$$

It follows that a bundle morphism (Φ, ϕ) yields a morphism of simple graded manifolds

$$(Z, \mathfrak{A}_E) \to (Z', \mathfrak{A}_{E'}). \tag{30}$$

This is a pair $(\phi, \widehat{\Phi} = \phi_* \circ \Phi^*)$ of a morphism ϕ of body manifolds and the composition $\phi_* \circ \Phi^*$ of the pull-back

$$\mathcal{A}_{E'} \ni f \to \Phi^* f \in \mathcal{A}_E$$

of graded functions and the direct image ϕ_* of a sheaf $\mathfrak{A}_E$ onto Z'. Relative to local bases (z^A, c^a) and (z'^A, c'^a) for $(Z, \mathfrak{A}_E)$ and $(Z', \mathfrak{A}_{E'})$, the morphism (30) of simple graded manifolds reads

$$z' = \phi(z), \qquad \widehat{\Phi}(c'^a) = \Phi^a_b(z) c^b.$$

For instance, the graded manifold morphism (30) is a monomorphism (resp. epimorphism) if Φ is a bundle injection (resp. surjection).

In particular, let $(Y, \mathfrak{A})$ be a graded manifold whose body $Z = Y$ is a fibre bundle $\pi : Y \to X$. Let us consider the trivial graded manifold $(X, \mathfrak{A}_0 = C^\infty_X)$ (21). Then we have a graded bundle

$$(Y, \mathfrak{A}) \to (X, C^\infty_X) \tag{31}$$

in accordance with Definition 3.

Definition 4. We agree to call the graded bundle (31) over a trivial graded manifold (X, C^∞_X) the graded bundle over a smooth manifold. Let us denote it by $(X, Y, \mathfrak{A})$.

If $Y \to X$ is a vector bundle, this is a particular case of graded fibre bundles in [16, 27] when their base is a trivial graded manifold.

Remark 2. Let $Y \to X$ be a fibre bundle. Then a trivial graded manifold (Y, C^∞_Y) together with a real ring monomorphism $C^\infty(X) \to C^\infty(Y)$ is the graded bundle (X, Y, C^∞_Y) (31).

Remark 3. A graded manifold $(X, \mathfrak{A})$ itself can be treated as the graded bundle $(X, X, \mathfrak{A})$ (31) associated to the identity smooth bundle $X \to X$.

Given a graded bundle $(X, Y, \mathfrak{A})$, a local basis for a graded manifold $(Y, \mathfrak{A})$ can be brought into a form (x^λ, y^i, c^a) where (x^λ, y^i) are bundle coordinates of $Y \to X$.

Let $(Y, \mathfrak{A}_F)$ be a simple graded manifold modelled over a vector bundle $F \to Y$. This is a graded bundle $(X, Y, \mathfrak{A}_F)$ together with a composite bundle

$$F \to Y \to X. \tag{32}$$

Accordingly, there is a composite bundle

$$\bigwedge_Y F^* \to Y \to X$$

so that the structure ring $\mathcal{A}_F = \wedge F^*(Y)$ (20) of graded functions on a simple graded manifold $(Y, \mathfrak{A}_F)$ is a graded commutative $C^\infty(X)$-ring. Let the composite bundle (32) be provided with adapted bundle coordinates (x^λ, y^i, q^a) possessing transition functions

$$x'^\lambda(x^\mu), \qquad y'^i(x^\mu, y^j), \qquad q'^a = \rho^a_b(x^\mu, y^j)q^b.$$

Then the corresponding local basis for a graded manifold is (x^λ, y^i, c^a) together with transition functions

$$x'^\lambda(x^\mu), \qquad y'^i(x^\mu, y^j), \qquad c'^a = \rho^a_b(x^\mu, j^j)c^b.$$

We call it the local basis for a graded bundle $(X, Y, \mathfrak{A}_F)$.

Any global section s of a fibre bundle $Y \to X$ yields the pull-back vector bundle $s^*F \to X$ over X which is a subbundle of a fibre bundle $Y \to X$ [31, 33]. Let $(X, \mathfrak{A}_{s^*F})$ be a simple graded manifold modelled over the pull-back bundle s^F. Its structure ring consists of global sections of the pull-back bundle

$$s^* \wedge F^* = \wedge s^* F^* \to X, \tag{33}$$

which is a subbundle of a fibre bundle $\wedge F^* \to X$. With a global section s of a fibre bundle $Y \to X$, any global section f of an exterior bundle $\wedge F^* \to Y$ defines a global section $f \circ s$ of the exterior bundle (33). The correspondence $f \to f \circ s$ yields an even morphism of a graded commutative $C^\infty(X)$-ring $\mathcal{A}_F$ to a graded commutative ring $\mathcal{A}_{s^*F}$. This morphism is an epimorphism because any global section of a subbundle $s^* \wedge F^* \subset \wedge F^* \to X$ is a local section of $\wedge F^* \to Y$ over a

closed subset $s(X) \subset Y$ and, therefore, it can be extended to a global section f of $\wedge F^* \to Y$. Passing to the sheafs $\mathfrak{A}_F$ and $\mathfrak{A}_{s^*F}$ of germs of graded functions, we come to the following.

Theorem 5. Given a graded bundle $(X, Y, \mathfrak{A}_F)$, any global section $s : X \to Y$ of a fibre bundle $Y \to X$ yields a monomorphism of a graded manifold $(X, \mathfrak{A}_{s^*F})$ to a graded manifold $(Y, \mathfrak{A}_F)$.

6 Graded jet manifolds

As was mentioned above, Lagrangian theory of even fields represented by sections of a fibre bundle $Y \to X$ is formulated on jet manifolds J^*Y of $Y \to X$. Therefore, one must define jets of odd fields.

Given a graded manifold $(X, \mathfrak{A})$ and its structure ring $\mathcal{A}$, one can define the jet module $J^1\mathcal{A}$ of a $C^\infty(X)$-ring $\mathcal{A}$ [17, 32]. If $(X, \mathfrak{A}_E)$ is a simple graded manifold modelled over a vector bundle $E \to X$, the jet module $J^1\mathcal{A}_E$ is a module of global sections of the jet bundle $J^1(\wedge E^*)$. From the mathematical viewpoint, a problem is that $J^1\mathcal{A}_E$ fails to be a structure ring of some graded manifold. From the physical viewpoint, elements of $J^1\mathcal{A}_E$ are affine in jets of odd fields.

By this reason, we have suggested a different construction of jets of graded manifolds, though it is applied only to simple graded manifolds [20, 33, 34].

Let $(X, \mathcal{A}_E)$ be a simple graded manifold modelled over a vector bundle $E \to X$. Let us consider a k-order jet manifold J^kE of E It is a vector bundle over X. Then let $(X, \mathcal{A}_{J^kE})$ be a simple graded manifold modelled over $J^kE \to X$.

Definition 6. We agree to call $(X, \mathcal{A}_{J^kE})$ the graded k-order jet manifold of a simple graded manifold $(X, \mathcal{A}_E)$.

Given a splitting domain $(U; x^\lambda, c^a)$ of a graded manifold $(Z, \mathcal{A}_E)$, we have a splitting domain

$$(U; x^\lambda, c^a, c^a_\lambda, c^a_{\lambda_1\lambda_2}, \dots c^a_{\lambda_1\dots\lambda_k}), \qquad c'^a_{\lambda\lambda_1\dots\lambda_r} = \rho^a_b(x)c^a_{\lambda\lambda_1\dots\lambda_r} + \partial_\lambda\rho^a_b(x)c^a_{\lambda_1\dots\lambda_r},$$

of a graded jet manifold $(X, \mathcal{A}_{J^kE})$.

Remark 4. Given a manifold X and its coordinate chart $(U; x^\lambda)$, a multi-index Λ of the length $|\Lambda| = k$ throughout denotes a collection of indices $(\lambda_1 \dots \lambda_k)$ modulo permutations. By $\lambda+\Lambda$ is meant a multi-index $(\lambda\lambda_1 \dots \lambda_k)$. Summation over a multi-index Λ means separate summation over each its index λ_i. We use the compact notation $\partial_\Lambda = \partial_{\lambda_k} \circ \dots \circ \partial_{\lambda_1}$ and $\Lambda = (\lambda_1 \dots \lambda_k)$.

As was mentioned above, a graded manifold is a particular graded bundle over its body (Remark 3). Then Definition 6 of graded jet manifolds is generalized to graded bundles over smooth manifolds as follows.

Let $(X, Y, \mathfrak{A}_F)$ be a graded bundle modelled over the composite bundle (32). It is readily observed that the jet manifold J^rF of $F \to X$ is a vector bundle $J^rF \to J^rY$ coordinated by $(x^\lambda, y^i_\Lambda, q^a_\Lambda)$, $0 \leq |\Lambda| \leq r$. Let $(J^rY, \mathfrak{A}_r = \mathfrak{A}_{J^rF})$ be a simple graded manifold modelled over this vector bundle. Its local basis is $(x^\lambda, y^i_\Lambda, c^a_\Lambda)$, $0 \leq |\Lambda| \leq r$.

Definition 7. We call $(J^rY, \mathfrak{A}_r)$ the graded r-order jet manifold of a graded bundle $(X, Y, \mathfrak{A}_F)$.

In particular, let $Y \to X$ be a smooth bundle seen as a trivial graded bundle (X, Y, C^∞_Y) modelled over a composite bundle $Y \times 0 \to Y \to X$. Then its graded jet manifold in accordance with Definition 7 is a trivial graded bundle $(X, J^rY, C^\infty_{J^rY})$, i.e., a jet manifold J^rY of Y.

Thus, Definition 7 of graded jet manifolds of graded bundles is compatible with the conventional definition of jets of fibre bundles. As was mentioned above, it however differs from that of jet graded bundles in [16, 27].

Jet manifolds J^*Y of a fibre bundle $Y \to X$ form the inverse sequence (2) where π^r_{r-1} are affine bundles. One can think of elements of its projective limit $J^\infty Y$ as being infinite order jets of sections of $Y \to X$ identified by their Taylor series at points of X. The set $J^\infty Y$ is endowed with the projective limit topology which makes $J^\infty Y$ into a paracompact Fréchet manifold [20, 37]. It is called the infinite order jet manifold and is provided with with the adapted coordinates

$$(x^\lambda, y^i_\Lambda), \quad 0 \leq |\Lambda|, \qquad y'^i_{\lambda+\Lambda} = \frac{\partial x^\mu}{\partial x'^\lambda} d_\mu y'^i_\Lambda, \tag{34}$$

where

$$d_\lambda = \partial_\lambda + y^i_\lambda \partial_i + \sum_{0<|\Lambda|} y^i_{\lambda+\Lambda} \partial^\Lambda_i$$

are total derivatives. The inverse sequence (2) of jet manifolds yields the direct sequence of graded differential algebras $\mathcal{O}^*_r$ of exterior forms on finite order jet manifolds

$$\mathcal{O}^*(X) \xrightarrow{\pi^*} \mathcal{O}^*(Y) \xrightarrow{\pi^{1*}_0} \mathcal{O}^*_1 \longrightarrow \cdots \mathcal{O}^*_{r-1} \xrightarrow{\pi^{r}_{r-1}{}^*} \mathcal{O}^*_r \longrightarrow \cdots, \tag{35}$$

where $\pi^r_{r-1}{}^*$ are the pull-back monomorphisms. Its direct limit $\mathcal{O}^*_\infty = \varinjlim \mathcal{O}^*_r$ exists and consists of all exterior forms on finite order jet manifolds modulo the pull-back identification. The $\mathcal{O}^*_\infty$ is a differential graded algebra which inherits the operations of the exterior differential d and exterior product $\wedge$ of exterior algebras $\mathcal{O}^*_r$. A key point is that the cohomology $H^*(\mathcal{O}^*_\infty)$ of the de Rham complex

$$0 \longrightarrow \mathbb{R} \longrightarrow \mathcal{O}^0_\infty \xrightarrow{d} \mathcal{O}^1_\infty \xrightarrow{d} \cdots \tag{36}$$

of a differential graded algebra $\mathcal{O}^*_\infty$ equals the de Rham cohomology of a fibre bundle Y [2, 20].

Surjections $\pi^{r+1}_r : J^{r+1}Y \to J^rY$ (2) yield epimorphisms of graded manifolds

$$(\pi^{r+1}_r, \widehat{\pi}^{r+1}_r) : (J^{r+1}Y, \mathfrak{A}_{r+1}) \to (J^rY, \mathfrak{A}_r),$$

including the sheaf monomorphisms

$$\widehat{\pi}^{r+1}_r : \pi^{r+1*}_r \mathfrak{A}_r \to \mathfrak{A}_{r+1}, \tag{37}$$

where $\pi^{r+1*}_r \mathfrak{A}_r$ is the pull-back onto $J^{r+1}Y$ of a continuous fibre bundle $\mathfrak{A}_r \to J^rY$. As a consequence, we have the inverse sequence of graded manifolds (3).

The sheaf monomorphism (37) induces a monomorphism of canonical presheaves $\overline{\mathfrak{A}}_r \to \overline{\mathfrak{A}}_{r+1}$, which associates to each open subset $U \subset J^{r+1}Y$ the ring of sections of $\mathfrak{A}_r$ over $\pi^{r+1}_r(U)$. Accordingly, there is a monomorphism of the structure rings

$$\pi^{r+1*}_r : \mathcal{S}^0_r[F; Y] \to \mathcal{S}^0_{r+1}[F; Y] \tag{38}$$

of graded functions on graded manifolds $(J^rY, \mathfrak{A}_r)$ and $(J^{r+1}Y, \mathfrak{A}_{r+1})$. Since the differential calculus $\mathcal{S}^*_r[F;Y]$ and $\mathcal{S}^*_{r+1}[F;Y]$ are minimal. Therefore, the monomorphism (38) yields a monomorphism of differential bigraded algebras

$$\pi_r^{r+1*} : \mathcal{S}^*_r[F;Y] \to \mathcal{S}^*_{r+1}[F;Y]. \tag{39}$$

As a consequence, we have the direct system of differential bigraded algebras

$$\mathcal{S}^*[F;Y] \xrightarrow{\pi^*} \mathcal{S}^*_1[F;Y] \longrightarrow \cdots \mathcal{S}^*_{r-1}[F;Y] \xrightarrow{\pi^{r*}_{r-1}} \mathcal{S}^*_r[F;Y] \longrightarrow \cdots . \tag{40}$$

The differential bigraded algebra $\mathcal{S}^*_\infty[F;Y]$ that we associate to a graded bundle $(Y, \mathfrak{A}_F)$ is defined as the direct limit

$$\mathcal{S}^*_\infty[F;Y] = \varinjlim \mathcal{S}^*_r[F;Y] \tag{41}$$

of the direct system (40). It consists of all graded exterior forms $\phi \in \mathcal{S}^*[F_r; J^rY]$ on graded manifolds $(J^rY, \mathfrak{A}_r)$ modulo the monomorphisms (39). Its elements obey the relations (14).

The cochain monomorphisms $\mathcal{O}^*_r \to \mathcal{S}^*_r[F;Y]$ (28) provide a monomorphism of the direct system (35) to the direct system (40) and, consequently, a monomorphism $\mathcal{O}^*_\infty \to \mathcal{S}^*_\infty[F;Y]$ of their direct limits. In particular, $\mathcal{S}^*_\infty[F;Y]$ is an $\mathcal{O}^0_\infty$-algebra.

One can think of elements of $\mathcal{S}^*_\infty[F;Y]$ as being graded differential forms on an infinite order jet manifold $J^\infty Y$ [20]. In particular, one can restrict $\mathcal{S}^*_\infty[F;Y]$ to the coordinate chart (34) of $J^\infty Y$ and say that $\mathcal{S}^*_\infty[F;Y]$ as an $\mathcal{O}^0_\infty$-algebra is locally generated by the elements

$$(c^a_\Lambda, dx^\lambda, \theta^a_\Lambda = dc^a_\Lambda - c^a_{\lambda+\Lambda} dx^\lambda, \theta^i_\Lambda = dy^i_\Lambda - y^i_{\lambda+\Lambda} dx^\lambda), \qquad 0 \leq |\Lambda|,$$

where c^a_Λ, θ^a_Λ are odd and dx^λ, θ^i_Λ are even. We agree to call (y^i, c^a) the local generating basis for $\mathcal{S}^*_\infty[F;Y]$. Let the collective symbol s^A stand for its elements. Accordingly, the notation s^A_Λ and $\theta^A_\Lambda = ds^A_\Lambda - s^A_{\lambda+\Lambda} dx^\lambda$ is introduced. For the sake of simplicity, we further denote $[A] = [s^A]$.

7 Lagrangian theory of even and odd fields

A differential bigraded algebra $\mathcal{S}^*_\infty[F;Y]$ is decomposed into $\mathcal{S}^0_\infty[F;Y]$-modules $\mathcal{S}^{k,r}_\infty[F;Y]$ of k-contact and r-horizontal graded forms together with the corresponding projections

$$h_k : \mathcal{S}^*_\infty[F;Y] \to \mathcal{S}^{k,*}_\infty[F;Y], \qquad h^m : \mathcal{S}^*_\infty[F;Y] \to \mathcal{S}^{*,m}_\infty[F;Y].$$

Accordingly, the graded exterior differential d on $\mathcal{S}^*_\infty[F;Y]$ falls into a sum $d = d_V + d_H$ of the vertical graded differential

$$d_V \circ h^m = h^m \circ d \circ h^m, \qquad d_V(\phi) = \theta^A_\Lambda \wedge \partial^\Lambda_A \phi, \qquad \phi \in \mathcal{S}^*_\infty[F;Y],$$

and the total graded differential

$$d_H \circ h_k = h_k \circ d \circ h_k, \qquad d_H \circ h_0 = h_0 \circ d, \qquad d_H(\phi) = dx^\lambda \wedge d_\lambda(\phi),$$

where

$$d_\lambda = \partial_\lambda + \sum_{0 \leq |\Lambda|} s^A_{\lambda+\Lambda} \partial^\Lambda_A$$

are the graded total derivatives. These differentials obey the nilpotent relations

$$d_H \circ d_H = 0, \qquad d_V \circ d_V = 0, \qquad d_H \circ d_V + d_V \circ d_H = 0.$$

Similarly to a differential graded algebra $\mathcal{O}^*_\infty$, a differential bigraded algebra $\mathcal{S}^*_\infty[F;Y]$ is provided with the graded projection endomorphism

$$\begin{aligned}
&\varrho = \sum_{k>0} \frac{1}{k} \overline{\varrho} \circ h_k \circ h^n : \mathcal{S}^{*>0,n}_\infty[F;Y] \to \mathcal{S}^{*>0,n}_\infty[F;Y],\\
&\overline{\varrho}(\phi) = \sum_{0\leq|\Lambda|} (-1)^{|\Lambda|}\theta^A \wedge [d_\Lambda(\partial^\Lambda_A \rfloor \phi)], \qquad \phi \in \mathcal{S}^{>0,n}_\infty[F;Y],
\end{aligned}$$

such that $\varrho \circ d_H = 0$, and with the nilpotent graded variational operator

$$\delta = \varrho \circ d\mathcal{S}^{*,n}_\infty[F;Y] \to \mathcal{S}^{*+1,n}_\infty[F;Y]. \tag{42}$$

With these operators a differential bigraded algebra $\mathcal{S}^*_\infty[F;Y]$ is decomposed into the Grassmann-graded variational bicomplex [20, 29, 34].

We restrict our consideration to its short variational subcomplex

$$\begin{aligned}
&0 \to \mathbb{R} \to \mathcal{S}^0_\infty[F;Y] \xrightarrow{d_H} \mathcal{S}^{0,1}_\infty[F;Y] \cdots \xrightarrow{d_H} \mathcal{S}^{0,n}_\infty[F;Y] \xrightarrow{\delta} \mathbf{E}_1, \qquad (43)\\
&\mathbf{E}_1 = \varrho(\mathcal{S}^{1,n}_\infty[F;Y]), \qquad n = \dim X,
\end{aligned}$$

and the subcomplex of one-contact graded forms

$$0 \to \mathcal{S}^{1,0}_\infty[F;Y] \xrightarrow{d_H} \mathcal{S}^{1,1}_\infty[F;Y] \cdots \xrightarrow{d_H} \mathcal{S}^{1,n}_\infty[F;Y] \xrightarrow{\varrho} \mathbf{E}_1 \to 0. \tag{44}$$

They possess the following cohomology [18, 29, 34].

Theorem 8. Cohomology of the complex (43) equals the de Rham cohomology $H^*_{DR}(Y)$ of Y. The complex (44) is exact.

Decomposed into a variational bicomplex, the differential bigraded algebra $\mathcal{S}^*_\infty[F;Y]$ describes Grassmann-graded field theory on a graded bundle $(X,Y,\mathfrak{A}_F)$. Its graded Lagrangian is defined as an element

$$L = \mathcal{L}\omega \in \mathcal{S}^{0,n}_\infty[F;Y]$$

of the graded variational complex (43). Accordingly, a graded exterior form

$$\delta L = \theta^A \wedge \mathcal{E}_A\omega = \sum_{0\leq|\Lambda|} (-1)^{|\Lambda|}\theta^A \wedge d_\Lambda(\partial^\Lambda_A L)\omega \in \mathbf{E}_1 \tag{45}$$

is said to be its graded Euler–Lagrange operator. We agree to call a pair $(\mathcal{S}^{0,n}_\infty[F;Y], L)$ the Grassmann-graded (or, simply, graded) Lagrangian system and $\mathcal{S}^*_\infty[F;Y]$ the field system algebra.

The following is a corollary of Theorem 8 [20, 34].

Theorem 9. Every d_H-closed graded form $\phi \in \mathcal{S}^{0,m<n}_\infty[F;Y]$ falls into the sum

$$\phi = h_0\sigma + d_H\xi, \qquad \xi \in \mathcal{S}^{0,m-1}_\infty[F;Y], \tag{46}$$

where σ is a closed m-form on Y. Any δ-closed (i.e., variationally trivial) graded Lagrangian $L \in \mathcal{S}^{0,n}_\infty[F;Y]$ is a sum

$$L = h_0\sigma + d_H\xi, \qquad \xi \in \mathcal{S}^{0,n-1}_\infty[F;Y], \tag{47}$$

where σ is a closed n-form on Y.

Corollary 10. Any variationally trivial odd Lagrangian is d_H-exact.

The exactness of the complex (44) at the term $\mathcal{S}^{1,n}_\infty[F;Y]$ results in the following [18, 20, 34].

Theorem 11. Given a graded Lagrangian L, there is the decomposition

$$dL = \delta L - d_H\Xi_L, \qquad \Xi \in \mathcal{S}^{n-1}_\infty[F;Y], \tag{48}$$

$$\Xi_L = L + \sum_{s=0} \theta^A_{\nu_s\ldots\nu_1} \wedge F^{\lambda\nu_s\ldots\nu_1}_A \omega_\lambda, \tag{49}$$

$$F^{\nu_k\ldots\nu_1}_A = \partial^{\nu_k\ldots\nu_1}_A\mathcal{L} - d_\lambda F^{\lambda\nu_k\ldots\nu_1}_A + \sigma^{\nu_k\ldots\nu_1}_A, \qquad k = 1,2,\ldots,$$

where local graded functions σ obey the relations

$$\sigma^\nu_A = 0, \qquad \sigma^{(\nu_k\nu_{k-1})\ldots\nu_1}_A = 0.$$

The form Ξ_L (49) provides a global Lepage equivalent of a graded Lagrangian L. In particular, one can locally choose Ξ_L (49) where all functions σ vanish.

Given a graded Lagrangian system $(\mathcal{S}^*_\infty[F;Y], L)$, by its infinitesimal transformations are meant contact graded derivations of the real graded commutative ring $\mathcal{S}^0_\infty[F;Y]$. They constitute a $\mathcal{S}^0_\infty[F;Y]$-module $\mathfrak{d}\mathcal{S}^0_\infty[F;Y]$ which is a real Lie superalgebra with respect to the Lie superbracket (10).

Theorem 12. The derivation module $\mathfrak{d}\mathcal{S}^0_\infty[F;Y]$ is isomorphic to the $\mathcal{S}^0_\infty[F;Y]$-dual $(\mathcal{S}^1_\infty[F;Y])^*$ of the module of graded one-forms $\mathcal{S}^1_\infty[F;Y]$. It follows that the differential bigraded algebra $\mathcal{S}^*_\infty[F;Y]$ is minimal differential calculus over the real graded commutative ring $\mathcal{S}^0_\infty[F;Y]$ [20, 34].

Let $\vartheta\rfloor\phi$, $\vartheta \in \mathfrak{d}\mathcal{S}^0_\infty[F;Y]$, $\phi \in \mathcal{S}^1_\infty[F;Y]$, denote the corresponding interior product. Extended to the differential bigraded algebra $\mathcal{S}^*_\infty[F;Y]$, it obeys the rule

$$\vartheta\rfloor(\phi\wedge\sigma) = (\vartheta\rfloor\phi)\wedge\sigma + (-1)^{|\phi|+[\phi][\vartheta]}\phi\wedge(\vartheta\rfloor\sigma), \qquad \phi,\sigma \in \mathcal{S}^*_\infty[F;Y].$$

Restricted to a coordinate chart (34) of $J^\infty Y$, the algebra $\mathcal{S}^*_\infty[F;Y]$ is a free $\mathcal{S}^0_\infty[F;Y]$-module generated by one-forms dx^λ, θ^A_Λ. Due to the isomorphism stated in Theorem 12, any graded derivation $\vartheta \in \mathfrak{d}\mathcal{S}^0_\infty[F;Y]$ takes a local form

$$\vartheta = \vartheta^\lambda\partial_\lambda + \vartheta^A\partial_A + \sum_{0<|\Lambda|} \vartheta^A_\Lambda\partial^\Lambda_A, \tag{50}$$

where $\partial^\Lambda_A\rfloor dy^B_\Sigma = \delta^B_A\delta^\Lambda_\Sigma$ up to permutations of multi-indices Λ and Σ.

Every graded derivation ϑ (50) yields the graded Lie derivative

$$\mathbf{L}_\vartheta\phi = \vartheta\rfloor d\phi + d(\vartheta\rfloor\phi), \qquad \phi \in \mathcal{S}^*_\infty[F;Y],$$
$$\mathbf{L}_\vartheta(\phi\wedge\sigma) = \mathbf{L}_\vartheta(\phi)\wedge\sigma + (-1)^{[\vartheta][\phi]}\phi\wedge\mathbf{L}_\vartheta(\sigma),$$

of the differential bigraded algebra $\mathcal{S}^*_\infty[F;Y]$. A graded derivation ϑ (50) is called contact if the Lie derivative $\mathbf{L}_\vartheta$ preserves the ideal of contact graded forms of the differential bigraded algebra $\mathcal{S}^*_\infty[F;Y]$.

With respect to the local generating basis (s^A) for the differential bigraded algebra $\mathcal{S}^*_\infty[F;Y]$, any its contact graded derivation takes a form

$$\vartheta = \upsilon_H + \upsilon_V = \upsilon^\lambda d_\lambda + [\upsilon^A\partial_A + \sum_{|\Lambda|>0} d_\Lambda(\upsilon^A - s^A_\mu\upsilon^\mu)\partial^\Lambda_A], \tag{51}$$

where υ_H and υ_V denotes the horizontal and vertical parts of ϑ. In particular, a vertical contact graded derivation reads

$$\vartheta = \upsilon^A\partial_A + \sum_{|\Lambda|>0} d_\Lambda\upsilon^A\partial^\Lambda_A. \tag{52}$$

A glance at the expression (51) shows that a contact graded derivation ϑ as an infinite order jet prolongation of its restriction

$$\upsilon = \upsilon^\lambda\partial_\lambda + \upsilon^A\partial_A \tag{53}$$

to the graded commutative ring $S^0[F;Y]$. We call υ (53) the generalized graded vector field.

A corollary of the decomposition (48) is the first variational formula for a graded Lagrangian [5, 18, 20].

Theorem 13. The Lie derivative of a graded Lagrangian along any contact graded derivation (51) obeys the first variational formula

$$\mathbf{L}_\vartheta L = \upsilon_V\rfloor\delta L + d_H(h_0(\vartheta\rfloor\Xi_L)) + d_V(\upsilon_H\rfloor\omega)\mathcal{L}, \tag{54}$$

where Ξ_L is the Lepage equivalent (49) of L.

A contact graded derivation ϑ (51) is called a variational symmetry of a graded Lagrangian L if the Lie derivative $\mathbf{L}_\vartheta L$ is d_H-exact, i.e., $\mathbf{L}_\vartheta L = d_H\sigma$. Then a corollary of the first variational formula (54) is the first Noether theorem for graded Lagrangians.

Theorem 14. If a contact graded derivation ϑ (51) is a variational symmetry of a graded Lagrangian L, the first variational formula (54) leads to the weak conservation law

$$0 \approx d_H(h_0(\vartheta\rfloor\Xi_L) - \sigma) \tag{55}$$

on the shell $\operatorname{Ker}\delta L$.

For the sake of brevity, the common symbol υ further stands for the generalized graded vector field υ (53), the contact graded derivation ϑ (51) determined by υ, and a Lie derivative $\mathbf{L}_\vartheta$.

A vertical contact graded derivation $\upsilon = \upsilon^A\partial_A$ is said to be nilpotent if $\upsilon(\upsilon\phi) = 0$ for any horizontal graded form $\phi \in S^{0,*}_\infty[F,Y]$. It is nilpotent only if it is odd and iff the equality $\upsilon(\upsilon^A) = 0$ holds for all υ^A [18].

Remark 5. For the sake of convenience, right derivations $\overleftarrow{\upsilon} = \overleftarrow{\partial}_A\upsilon^A$ also are considered. They act on graded functions and differential forms ϕ on the right by the rules

$$\overleftarrow{\upsilon}(\phi) = d\phi\lfloor\overleftarrow{\upsilon} + d(\phi\lfloor\overleftarrow{\upsilon}), \qquad \overleftarrow{\upsilon}(\phi\wedge\phi') = (-1)^{[\phi']}\overleftarrow{\upsilon}(\phi)\wedge\phi' + \phi\wedge\overleftarrow{\upsilon}(\phi').$$

8 Lagrangian BRST theory

As was mentioned above, quantization of Lagrangian field theory essentially depends on its degeneracy characterized by a hierarchy of Noether identities, and a first step to this quantization is BRST extension of an original Lagrangian theory [3, 20, 21].

Any Lagrangian system admits Noether identities. Therefore, a problem lies in separation of trivial and non-trivial Noether identities that one can perform in the homology terms so that trivial Noether identities are boundaries, whereas the non-trivial ones are cycles of some chain complex [6, 7, 20].

Without a lose of generality, let a Lagrangian L of a graded Lagrangian system $(\mathcal{S}^{0,n}_\infty[F;Y],L)$ be even. Its Euler–Lagrange Euler–Lagrange operator $\delta L\in\mathbf{E}_1\subset\mathcal{S}^{1,n}_\infty[F;Y]$ (45) takes the values into a vector bundle

$$\overline{VF}=V^*F\underset{F}{\otimes}\overset{n}{\wedge}T^*X\to F, \tag{56}$$

where V^*F is the vertical cotangent bundle of $F\to X$. For our purpose of studying gauge theory, it suffices to consider a pull-back composite bundle F (32) that is

$$F=Y\underset{X}{\times}F^1\to Y\to X, \tag{57}$$

where $F^1\to X$ is a vector bundle.

Let us introduce the following notation. Given the vertical tangent bundle VE of a fibre bundle $E\to X$, by its density-dual bundle is meant a fibre bundle

$$\overline{VE}=V^*E\underset{E}{\otimes}\overset{n}{\wedge}T^*X. \tag{58}$$

If $E\to X$ is a vector bundle, we have

$$\overline{VE}=\overline{E}\underset{X}{\times}E,\qquad \overline{E}=E^*\underset{X}{\otimes}\overset{n}{\wedge}T^*X,$$

where $\overline{E}$ is called the density-dual of E. Let $E=E^0\oplus_X E^1$ be a graded vector bundle over X. Its graded density-dual is defined to be $\overline{E}=\overline{E}^1\oplus_X\overline{E}^0$. In these terms, we treat the composite bundle F (32) as a graded vector bundle over Y possessing only an odd part. The density-dual $\overline{VF}$ (58) of the vertical tangent bundle VF of $F\to X$ is $\overline{VF}$ (56). If F (32) is the pull-back bundle (57), then

$$\overline{VF}=((\overline{F}^1\underset{Y}{\oplus}V^*Y)\underset{Y}{\otimes}\overset{n}{\wedge}T^*X)\underset{Y}{\oplus}F^1 \tag{59}$$

is a graded vector bundle over Y. Given a graded vector bundle $E=E^0\oplus_Y E^1$ over Y, we consider a composite bundle $E\to E^0\to X$ and the differential bigraded algebra (41):

$$\mathcal{P}^*_\infty[E;Y]=\mathcal{S}^*_\infty[E;E^0]. \tag{60}$$

Turn now to Noether identities of a Lagrangian system. One can associate to any graded Lagrangian system $(\mathcal{S}^*_\infty[F;Y],L)$ the chain complex (61) whose one-boundaries vanish on-shell as follows.

Let us consider the density-dual $\overline{VF}$ (59) of the vertical tangent bundle $VF\to F$, and let us enlarge the original field system algebra $\mathcal{S}^*_\infty[F;Y]$ to the differential bigraded algebra $\mathcal{P}^*_\infty[\overline{VF};Y]$ (60) with a local generating basis

$$(s^A,\overline{s}_A),\qquad [\overline{s}_A]=([A]+1)\mathrm{mod}\,2.$$

Following the conventional terminology of Lagrangian BRST theory [3, 21], we agree to call its elements $\overline{s}_A$ the antifields of antifield number $\mathrm{Ant}[\overline{s}_A] = 1$. The differential bigraded algebra $\mathcal{P}^*_\infty[\overline{VF}; Y]$ is endowed with the nilpotent right graded derivation

$$\overline{\delta} = \overleftarrow{\partial}{}^A \mathcal{E}_A,$$

where $\mathcal{E}_A$ are the variational derivatives (45). Then we have the chain complex

$$0 \leftarrow \mathrm{Im}\,\overline{\delta} \xleftarrow{\overline{\delta}} \mathcal{P}^{0,n}_\infty[\overline{VF}; Y]_1 \xleftarrow{\overline{\delta}} \mathcal{P}^{0,n}_\infty[\overline{VF}; Y]_2 \tag{61}$$

of graded densities of antifield number ≤ 2. Its one-boundaries $\overline{\delta}\Phi$, $\Phi \in \mathcal{P}^{0,n}_\infty[\overline{VF}; Y]_2$, by very definition, vanish on-shell.

Any one-cycle

$$\Phi = \sum_{0 \leq |\Lambda|} \Phi^{A,\Lambda} \overline{s}_{\Lambda A} \omega \in \mathcal{P}^{0,n}_\infty[\overline{VF}; Y]_1 \tag{62}$$

of the complex (61) is a differential operator on the bundle $\overline{VF}$ such that it is linear on fibres of $\overline{VF} \to F$ and its kernel contains the graded Euler–Lagrange operator δL (45), i.e.,

$$\overline{\delta}\Phi = 0, \qquad \sum_{0 \leq |\Lambda|} \Phi^{A,\Lambda} d_\Lambda \mathcal{E}_A \omega = 0. \tag{63}$$

Thus the one-cycles (62) define the Noether identities (63) of the Euler–Lagrange operator δL a graded Lagrangian system $(\mathcal{S}^*_\infty[F; Y], L)$.

In particular, one-chains Φ (62) are necessarily Noether identities if they are boundaries. Therefore, these Noether identities are called trivial. Accordingly, non-trivial Noether identities modulo the trivial ones are associated to elements of the first homology $H_1(\overline{\delta})$ of the complex (61). A Lagrangian system is called degenerate if there are non-trivial Noether identities.

Let us assume that a $C^\infty(X)$-module $H_1(\overline{\delta})$ is finitely generated. Namely, there exists a graded projective $C^\infty(X)$-module $\mathcal{C}_{(0)} \subset H_1(\overline{\delta})$ of finite rank possessing a local basis $\{\Delta_r \omega\}$:

$$\Delta_r \omega = \sum_{0 \leq |\Lambda|} \Delta^{A,\Lambda}_r \overline{s}_{\Lambda A} \omega, \qquad \Delta^{A,\Lambda}_r \in \mathcal{S}^0_\infty[F; Y], \tag{64}$$

such that any element $\Phi \in H_1(\overline{\delta})$ factorizes as

$$\Phi = \sum_{0 \leq |\Xi|} \Phi^{r,\Xi} d_\Xi \Delta_r \omega, \qquad \Phi^{r,\Xi} \in \mathcal{S}^0_\infty[F; Y], \tag{65}$$

through elements (64) of $\mathcal{C}_{(0)}$. Thus, all non-trivial Noether identities (63) result from the Noether identities

$$\overline{\delta}\Delta_r = \sum_{0 \leq |\Lambda|} \Delta^{A,\Lambda}_r d_\Lambda \mathcal{E}_A = 0, \tag{66}$$

called the complete Noether identities.

Hereafter, we restrict our consideration to Lagrangian systems whose non-trivial Noether identities are finitely generated because this is just the case of gauge theory (Section 9).

Lemma 15. If the homology $H_1(\overline{\delta})$ of the complex (61) is finitely generated in the above mentioned sense, this complex can be extended to the one-exact chain complex (69) with a

boundary operator whose nilpotency conditions are equivalent to the complete Noether identities (66).

Indeed, by virtue of Serre–Swan Theorem 2, the graded module $\mathcal{C}_{(0)}$ is isomorphic to a module of sections of the density-dual $\overline{E}_0$ of some graded vector bundle $E_0 \to X$. Let us enlarge $\mathcal{P}^*_\infty[\overline{VF}; Y]$ to the differential bigraded algebra

$$\overline{\mathcal{P}}^*_\infty\{0\} = \mathcal{P}^*_\infty[\overline{VF} \underset{Y}{\oplus} \overline{E}_0; Y] \tag{67}$$

possessing the local generating basis $(s^A, \overline{s}_A, \overline{c}_r)$ where $\overline{c}_r$ are Noether antifields of Grassmann parity $[\overline{c}_r] = ([\Delta_r] + 1) \mathrm{mod}\, 2$ and antifield number $\mathrm{Ant}[\overline{c}_r] = 2$. The differential bigraded algebra (67) is provided with an odd right graded derivation

$$\delta_0 = \overline{\delta} + \sum_{0 \leq |\Lambda|} \overleftarrow{\partial}{}^r \Delta^{A,\Lambda}_r \overline{s}_{\Lambda A}, \tag{68}$$

called the Koszul–Tate operator. It is nilpotent iff the complete Noether identities (66) hold.

Then δ_0 (68) is a boundary operator of a short chain complex, called the Koszul–Tate complex,

$$0 \leftarrow \mathrm{Im}\, \overline{\delta} \xleftarrow{\overline{\delta}} \mathcal{P}^{0,n}_\infty[\overline{VF}; Y]_1 \xleftarrow{\delta_0} \overline{\mathcal{P}}^{0,n}_\infty\{0\}_2 \xleftarrow{\delta_0} \overline{\mathcal{P}}^{0,n}_\infty\{0\}_3 \tag{69}$$

of graded densities of antifield number ≤ 3. Let $H_*(\delta_0)$ denote its homology. It is readily observed that $H_0(\delta_0) = H_1(\delta_1) = 0$. Let us consider the second homology $H_2(\delta_0)$ of the complex (69). Its two-chains read

$$\Phi = G + H = \sum_{0 \leq |\Lambda|} G^{r,\Lambda} \overline{c}_{\Lambda r} \omega + \sum_{0 \leq |\Lambda|, |\Sigma|} H^{(A,\Lambda)(B,\Sigma)} \overline{s}_{\Lambda A} \overline{s}_{\Sigma B} \omega. \tag{70}$$

Its two-cycles define the first-stage Noether identities

$$\delta_0 \Phi = 0, \qquad \sum_{0 \leq |\Lambda|} G^{r,\Lambda} d_\Lambda \Delta_r \omega = -\overline{\delta} H.$$

They are trivial either if a two-cycle Φ (70) is a δ_0-boundary or its summand G vanishes on-shell. A field theory is said to be irreducible if first-stage Noether identities are trivial. As was mentioned above, we restrict our consideration to this case.

The inverse second Noether theorem (Theorem 16), that we formulate in homology terms, associates to the Koszul–Tate complex (69) of non-trivial Noether identities the cochain sequence (73) with the ascent operator u (74) whose components are non-trivial gauge symmetries of Lagrangian system.

Let us start with the following notation. Given the differential bigraded algebra $\overline{\mathcal{P}}^*_\infty\{0\}$ (67), we consider a differential bigraded algebra

$$\mathcal{P}^*_\infty\{0\} = \mathcal{P}^*_\infty[F \underset{Y}{\oplus} E_0; Y], \tag{71}$$

possessing a local generating basis (s^A, c^r), $[c^r] = ([\overline{c}_r] + 1) \mathrm{mod}\, 2$, and a differential bigraded algebra

$$P^*_\infty\{0\} = \mathcal{P}^*_\infty[\overline{VF} \underset{Y}{\oplus} E_0 \underset{Y}{\oplus} \overline{E}_0; Y] \tag{72}$$

with a local generating basis $(s^A, \overline{s}_A, c^r, \overline{c}_r)$. Their elements c^r are called the ghosts of ghost number $\mathrm{gh}[c^r] = 1$ and antifield number $\mathrm{Ant}[c^r] = -1$. A $C^\infty(X)$-module $\mathcal{C}^{(0)}$ of ghosts is the

density-dual of a module $\mathcal{C}_{(0)}$ of antifields. The differential bigraded algebras $\overline{\mathcal{P}}^*_\infty\{0\}$ (67) and $\mathcal{P}^*_\infty\{0\}$ (71) are subalgebras of $P^*_\infty\{0\}$ (72). The operator δ_0 (68) is naturally extended to a graded derivation of the differential bigraded algebra $P^*_\infty\{0\}$.

Theorem 16. Given the Koszul–Tate complex (69), the module of graded densities $\mathcal{P}^{0,n}_\infty\{0\}$ contains a short cochain sequence

$$0 \to \mathcal{S}^{0,n}_\infty[F;Y] \xrightarrow{u} \mathcal{P}^{0,n}_\infty\{N\}^1, \tag{73}$$

$$u = u^A\frac{\partial}{\partial s^A}, \qquad u^A = \sum_{0\leq|\Lambda|} c^r_\Lambda \eta(\Delta^A_r)^\Lambda, \tag{74}$$

of ghost number ≤ 1. Its ascent operator u (74) is an odd graded derivation of ghost number 1 is a variational symmetry of a graded Lagrangian L.

A glance at the expression (74) shows that the variational symmetry u is a linear differential operator on the $C^\infty(X)$-module $\mathcal{C}^{(0)}$ of ghosts with values into the real space of variational symmetries. Therefore, u (74) is a gauge symmetry of a graded Lagrangian L which is associated to the complete Noether identities (66) [20, 30]. It is called the gauge operator.

Being a variational symmetry, the gauge operator u (74) defines the weak conservation law (55). Let u be an exact Lagrangian symmetry. In this case, the associated symmetry current

$$\mathcal{J}_u = -h_0(u\rfloor\Xi_L) \tag{75}$$

is conserved. The peculiarity of gauge conservation laws is that the symmetry current (75) is reduced to a superpotential as follows [20, 30].

Theorem 17. If u (74) is an exact gauge symmetry of a graded Lagrangian L, the corresponding conserved symmetry current $\mathcal{J}_u$ (75) takes a form

$$\mathcal{J}_u = W + d_H U = (W^\mu + d_\nu U^{\nu\mu})\omega_\mu, \tag{76}$$

where the term W vanishes on-shell, i.e., $W \approx 0$, and U is a horizontal graded $(n-2)$-form.

In contrast with the Koszul–Tate operator δ_0 (68), the gauge operator u (74) need not be nilpotent. Let us study its extension to a nilpotent graded derivation

$$\mathbf{b} = u + \gamma = u + \gamma^r\frac{\partial}{\partial c^r} \tag{77}$$

of ghost number 1 by means of antifield-free terms γ of higher polynomial degree in ghosts c^r and their jets c^r_Λ. We call $\mathbf{b}$ (77) the BRST operator, where gauge symmetries are extended to BRST transformations acting on ghosts [19, 20].

We refer to [20] for necessary conditions of the existence of such a BRST extension. If such an extension exists, a Lagrangian L also admits a certain BRST extension as follows.

The differential bigraded algebra $P^*_\infty\{0\}$ (72) is a particular field-antifield theory of the following type [3, 7, 21]. Let us consider a pull-back composite bundle

$$W = Z \underset{X}{\times} Z' \to Z \to X$$

where $Z' \to X$ is a vector bundle. Let us regard it as a graded vector bundle over Z possessing only odd part. The density-dual $\overline{VW}$ of the vertical tangent bundle VW of $W \to X$ is a graded

vector bundle

$$\overline{VW} = ((\overline{Z}' \underset{Z}{\oplus} V^*Z) \underset{Z}{\otimes} \overset{n}{\wedge} T^*X) \underset{Y}{\oplus} Z'$$

over Z (cf. (59)). Let us consider the differential bigraded algebra $\mathcal{P}^*_\infty[\overline{VW}; Z]$ (60) with the local generating basis

$$(z^a, \overline{z}_a), \qquad [\overline{z}_a] = ([z^a] + 1) \mathrm{mod}\, 2.$$

Its elements z^a and $\overline{z}_a$ are called fields and antifields, respectively.

Graded densities of this differential bigraded algebra are endowed with the antibracket

$$\{\mathfrak{L}\omega, \mathfrak{L}'\omega\} = \left[\frac{\overleftarrow{\delta}\mathfrak{L}}{\delta\overline{z}_a}\frac{\delta\mathfrak{L}'}{\delta z^a} + (-1)^{[\mathfrak{L}'](([\mathfrak{L}']+1)}\frac{\overleftarrow{\delta}\mathfrak{L}'}{\delta\overline{z}_a}\frac{\delta\mathfrak{L}}{\delta z^a}\right]\omega. \tag{78}$$

With this antibracket, one associates to any even Lagrangian $\mathfrak{L}\omega$ the odd vertical graded derivations

$$\upsilon_\mathfrak{L} = \overleftarrow{\mathcal{E}}{}^a\partial_a = \frac{\overleftarrow{\delta}\mathfrak{L}}{\delta\overline{z}_a}\frac{\partial}{\partial z^a}, \tag{79}$$

$$\overline{\upsilon}_\mathfrak{L} = \overleftarrow{\partial}{}^a\mathcal{E}_a = \frac{\overleftarrow{\partial}}{\partial\overline{z}_a}\frac{\delta\mathfrak{L}}{\delta z^a}, \tag{80}$$

$$\vartheta_\mathfrak{L} = \upsilon_\mathfrak{L} + \overline{\upsilon}^l_\mathfrak{L} = (-1)^{[a]+1}\left(\frac{\delta\mathfrak{L}}{\delta\overline{z}^a}\frac{\partial}{\partial z_a} + \frac{\delta\mathfrak{L}}{\delta z^a}\frac{\partial}{\partial\overline{z}_a}\right), \tag{81}$$

such that $\vartheta_\mathfrak{L}(\mathfrak{L}'\omega) = \{\mathfrak{L}\omega, \mathfrak{L}'\omega\}$.

Theorem 18. The following conditions are equivalent [20].

(i) The antibracket of a Lagrangian $\mathfrak{L}\omega$ is d_H-exact, i.e.,

$$\{\mathfrak{L}\omega, \mathfrak{L}\omega\} = 2\frac{\overleftarrow{\delta}\mathfrak{L}}{\delta\overline{z}_a}\frac{\delta\mathfrak{L}}{\delta z^a}\omega = d_H\sigma. \tag{82}$$

(ii) The graded derivations υ (79) and $\overline{\upsilon}$ (80) are variational symmetries of a Lagrangian $\mathfrak{L}\omega$.

(iii) The graded derivation $\vartheta_\mathfrak{L}$ (81) is nilpotent.

The equality (82) is called the classical master equation. A solution of the master equation (82) is called non-trivial if both the derivations (79) and (80) do not vanish. Being an element of the differential bigraded algebra $P^*_\infty\{0\}$ (72), an original Lagrangian L obeys the master equation (82) and yields the graded derivations $\upsilon_L = 0$ (79) and $\overline{\upsilon}_L = \overline{\delta}$ (80), i.e., it is a trivial solution of the master equation. One can show the following [20].

Theorem 19. The master equation has a non-trivial solution

$$L_E = L + \mathbf{b}(c^r\overline{c}_r) \tag{83}$$

if the gauge operator u (74) can be extended to the BRST operator $\mathbf{b}$ (77),

The non-trivial solution L_E (83) of the master equation is called the BRST extension of an original Lagrangian L. As was mentioned above, it is a necessary step towards quantization of classical Lagrangian field theory in terms of functional integrals.

9 Yang–Mills gauge theory

Yang–Mills gauge theory is a theory of principal connections on principal bundles [20, 34].

Let

$$\pi_P : P \to X \tag{84}$$

be a smooth principal bundle with a structure Lie group G which acts freely and transitively on P on the right by a fibrewise morphism

$$R_{GP} : G \underset{X}{\times} P \ni p \to pg \in P, \qquad \pi_P(p) = \pi_P(pg), \qquad p \in P. \tag{85}$$

As a consequence, the quotient of P with respect to the action (85) of G is diffeomorphic to a base X, i.e., $P/G = X$.

A principal G-bundle P is equipped with a bundle atlas

$$\Psi_P = \{(U_\alpha, \psi^P_\alpha), \varrho_{\alpha\beta}\} \tag{86}$$

whose trivialization morphisms

$$\psi^P_\alpha : \pi_P^{-1}(U_\alpha) \to U_\alpha \times G$$

obey the condition $\psi^P_\alpha(pg) = g\psi^P_\alpha(p)$, $g \in G$. Due to this property, every trivialization morphism ψ^P_α determines a unique local section $z_\alpha : U_\alpha \to P$ such that $(\psi^P_\alpha \circ z_\alpha)(x) = \mathbf{1}$, $x \in U_\alpha$. The transformation law for z_α reads

$$z_\beta(x) = z_\alpha(x)\varrho_{\alpha\beta}(x), \qquad x \in U_\alpha \cap U_\beta. \tag{87}$$

Conversely, the family $\{(U_\alpha, z_\alpha), \varrho_{\alpha\beta}\}$ of local sections of P which obey the transformation law (87) uniquely determines a bundle atlas Ψ_P of a principal bundle P.

In particular, it follows that a principal bundle admits a global section iff it is trivial.

Example 6. Let H be a closed subgroup of a real Lie group G. Then H is a Lie group. Let G/H be the quotient of G with respect to an action of H on G by right multiplications. Then

$$\pi_{GH} : G \to G/H \tag{88}$$

is a principal H-bundle. If H is a maximal compact subgroup of G, then G/H is diffeomorphic to $\mathbb{R}^m$ and, by virtue of the well known theorem, $G \to G/H$ is a trivial bundle, i.e., G is diffeomorphic to the product $\mathbb{R}^m \times H$.

Let us consider the tangent morphism

$$TR_{GP} : (G \times \mathfrak{g}_l) \underset{X}{\times} TP \underset{X}{\longrightarrow} TP \tag{89}$$

to the right action R_{GP} (85) of G on P. Its restriction to $T_{\mathbf{1}}G \times_X TP$ provides a homomorphism

$$\mathfrak{g}_l \ni \epsilon \to \xi_\epsilon \in \mathcal{T}(P) \tag{90}$$

of the left Lie algebra $\mathfrak{g}_l$ of G to the Lie algebra $\mathcal{T}(P)$ of vector fields on P. Vector fields ξ_ϵ (90) are obviously vertical. They are called fundamental vector fields [23]. Given a basis $\{\epsilon_r\}$ for $\mathfrak{g}_l$, the corresponding fundamental vector fields $\xi_r = \xi_{\epsilon_r}$ form a family of nowhere vanishing and

linearly independent sections of the vertical tangent bundle VP of $P \to X$. Consequently, this bundle is trivial

$$VP = P \times \mathfrak{g}_l. \tag{91}$$

Restricting the tangent morphism TR_{GP} (89) to

$$TR_{GP} : \widehat{0}(G) \underset{X}{\times} TP \underset{X}{\longrightarrow} TP, \tag{92}$$

we obtain the tangent prolongation of the structure group action R_{GP} (85). Since the action of G (85) on P is fibrewise, its action (92) is restricted to the vertical tangent bundle VP of P.

Taking the quotient of the tangent bundle $TP \to P$ and the vertical tangent bundle VP of P by G (92), we obtain the vector bundles

$$T_G P = TP/G, \qquad V_G P = VP/G \tag{93}$$

over X. Sections of $T_G P \to X$ are G-invariant vector fields on P. Accordingly, sections of $V_G P \to X$ are G-invariant vertical vector fields on P. Hence, a typical fibre of $V_G P \to X$ is the right Lie algebra $\mathfrak{g}_r$ of G subject to the adjoint representation of a structure group G. Therefore, $V_G P$ (93) is called the Lie algebra bundle.

Given a bundle atlas Ψ_P (86) of P, there is the corresponding atlas

$$\Psi = \{(U_\alpha, \psi_\alpha), \mathrm{Ad}_{\varrho_{\alpha\beta}}\} \tag{94}$$

of the Lie algebra bundle $V_G P$, which is provided with bundle coordinates $(U_\alpha; x^\mu, \chi^m)$ with respect to the fibre frames $\{e_m = \psi_\alpha^{-1}(x)(\varepsilon_m)\}$, where $\{\varepsilon_m\}$ is a basis for the Lie algebra $\mathfrak{g}_r$. These coordinates obey the transformation rule

$$\varrho(\chi^m)\varepsilon_m = \chi^m \mathrm{Ad}_{\varrho^{-1}}(\varepsilon_m).$$

A glance at this transformation rule shows that $V_G P$ is a bundle with a structure group G. Moreover, it is associated with a principal G-bundle P.

Accordingly, the vector bundle $T_G P$ (93) is endowed with bundle coordinates $(x^\mu, \dot{x}^\mu, \chi^m)$ with respect to the fibre frames $\{\partial_\mu, e_m\}$. Their transformation rule is

$$\varrho(\chi^m)\varepsilon_m = \chi^m \mathrm{Ad}_{\varrho^{-1}}(\varepsilon_m) + \dot{x}^\mu R^m_\mu \varepsilon_m.$$

The Lie bracket of G-invariant vertical vector fields on P goes to the quotient by G and defines the Lie bracket of sections of the vector bundle $T_V P \to X$. This bracket reads

$$[\xi, \eta] = c^r_{pq} \xi^p \eta^q e_r. \tag{95}$$

A glance at the expression (95) shows that sections of $V_G P$ form a finite-dimensional Lie $C^\infty(X)$-algebra, called the gauge algebra.

In classical gauge theory, gauge fields are conventionally described as principal connections on principal bundles. Principal connections on a principal bundle P (84) are connections on P which are equivariant with respect to the right action (85) of a structure group G on P. In order to describe them, we follow the definition of connections on a fibre bundle $Y \to X$ as global sections of the affine jet bundle $J^1 Y \to Y$ [20, 25, 34].

Let J^1P be the first order jet manifold of a principal G-bundle $P \to X$ (84). Then connections on a principal bundle $P \to X$ are global sections

$$A : P \to J^1P \tag{96}$$

of the affine jet bundle $J^1P \to P$. In order to define principal connections on $P \to X$, let us consider the jet prolongation

$$J^1R_{GP} : J^1(X \times G) \underset{X}{\times} J^1P \to J^1P$$

of the morphism R_{GP} (85). Restricting this morphism to

$$J^1R_G : \widehat{0}(G) \underset{X}{\times} J^1P \to J^1P,$$

we obtain the jet prolongation of the structure group action R_{GP} (85). It reads

$$G \ni g : j^1_x p \to (j^1_x p)g = j^1_x(pg). \tag{97}$$

Taking the quotient of the affine jet bundle J^1P by G (97), we obtain the affine bundle

$$C = J^1P/G \to X \tag{98}$$

modelled over the vector bundle

$$\overline{C} = T^*X \underset{X}{\otimes} V_GP \to X.$$

Hence, there is the vertical splitting $VC = C \otimes_X \overline{C}$ of the vertical tangent bundle VC of $C \to X$.

Remark 7. A glance at the expression (97) shows that the fibre bundle $J^1P \to C$ is a principal bundle with the structure group G. It is canonically isomorphic to the pull-back

$$J^1P = P_C = C \underset{X}{\times} P \to C. \tag{99}$$

Since a connection A (96) on a principal bundle $P \to X$ is principal connection if it is equivariant under the action (97) of a structure group G, there is obvious one-to-one correspondence between the principal connections on a principal G-bundle P and global sections

$$A : X \to C \tag{100}$$

of the factor bundle $C \to X$ (98), called the bundle of principal connections. Since the bundle $C \to X$ is affine, principal connections on a principal bundle always exist.

A bundle of principal connections C is provided with bundle coordinates (x^λ, a^m_μ) possessing a transformation rule

$$\varrho(a^m_\mu)\varepsilon_m = (a^m_\nu \mathrm{Ad}_{\varrho^{-1}}(\varepsilon_m) + R^m_\nu \varepsilon_m)\frac{\partial x^\nu}{\partial x'^\mu}. \tag{101}$$

Herewith, there is the canonical bundle monomorphism

$$\lambda_C : C \ni (x^\lambda, a^m_\mu) \underset{X}{\longrightarrow} dx^\mu \otimes (\partial_\mu + \chi^m_\mu e_m) \in T^*X \underset{X}{\otimes} T_GP.$$

Due to this bundle monomorphism, any principal connection A (100) is represented by a $T_G P$-valued form

$$A = dx^\lambda \otimes (\partial_\lambda + A^q_\lambda e_q) \tag{102}$$

which provides a splitting of this exact sequence

$$0 \to V_G P \underset{X}{\longrightarrow} T_G P \longrightarrow TX \to 0.$$

In classical gauge theory, global sections A (100) of the bundle $C \to X$ of principal connections are treated as gauge fields [20, 25, 34].

Let us consider first order Lagrangian theory of principal connections. Its configuration space is the first order jet manifold J^1C of the bundle of principal connections C (98), endowed with bundle coordinates (x^μ, a^m_μ) possessing transition functions (101). This configuration space admits the canonical splitting

$$J^1C = C_+ \underset{C}{\oplus} C_- = C_+ \underset{C}{\oplus} (C \underset{X}{\times} \overset{2}{\wedge} T^*X \otimes V_G P), \tag{103}$$

$$a^r_{\lambda\mu} = \frac{1}{2}(\mathcal{S}^r_{\lambda\mu} + \mathcal{F}^r_{\lambda\mu}) = \frac{1}{2}(a^r_{\lambda\mu} + a^r_{\mu\lambda} - c^r_{pq} a^p_\lambda a^q_\mu) + \frac{1}{2}(a^r_{\lambda\mu} - a^r_{\mu\lambda} + c^r_{pq} a^p_\lambda a^q_\mu). \tag{104}$$

Given a section A of a bundle $C \to X$ and its jet prolongation $j^1 A$, the projection of $j^1 A$ to C_- is the strength

$$\begin{aligned} F_A &= \frac{1}{2} F^r_{\lambda\mu} dx^\lambda \wedge dx^\mu \otimes e_r, \\ F^r_{\lambda\mu} &= \mathcal{F}^r_{\lambda\mu} \circ j^1 A = \partial_\lambda A^r_\mu - \partial_\mu A^r_\lambda + c^r_{pq} A^p_\lambda A^q_\mu, \end{aligned} \tag{105}$$

of a gauge field A.

Given a first order Lagrangian

$$L = \mathcal{L}\omega : J^1C \to \overset{n}{\wedge} T^*X \tag{106}$$

on J^1C, the corresponding Euler–Lagrange operator (45) reads

$$\mathcal{E}_L = \mathcal{E}^\mu_r \theta^r_\mu \wedge \omega = (\partial^\mu_r - d_\lambda \partial^{\lambda\mu}_r) \mathcal{L} \theta^r_\mu \wedge \omega. \tag{107}$$

Its kernel defines the Euler–Lagrange equation

$$\mathcal{E}^\mu_r = (\partial^\mu_r - d_\lambda \partial^{\lambda\mu}_r)\mathcal{L} = 0.$$

In Yang–Mills gauge theory, the Lagrangian L (106) is assumed to be invariant under principal gauge transformations. These are vertical principal automorphisms Φ_P of a principal G-bundle P which are equivariant under the right action (85) of a structure group G on P, i.e.,

$$\Phi_P(pg) = \Phi_P(p)g, \qquad g \in G, \qquad p \in P. \tag{108}$$

Every vertical principal automorphism of a principal bundle P is represented as $\Phi_P(p) = pf(p)$, $p \in P$, where f is a G-valued equivariant function on P, i.e.,

$$f(pg) = g^{-1} f(p) g, \qquad g \in G. \tag{109}$$

There is one-to-one correspondence between the equivariant functions f (109) and the global sections s of the P-associated group bundle $P^G \to X$ whose typical fibre is the group G which acts on itself by the adjoint representation. This correspondence is defined by the relation

$$s(\pi_P(p))p = pf(p), \qquad p \in P.$$

The group of vertical principal automorphisms of a principal G-bundle is called the gauge group. It is isomorphic to the group $P^G(X)$ of global sections of the group bundle P^G.

In order to describe principal gauge transformations of the Lagrangian L (106), let us restrict our consideration to (local) one-parameter groups of vertical principal automorphisms of P. Their infinitesimal generators are G-invariant vertical vector fields ξ on P. They are represented by sections

$$\xi = \xi^p e_p \tag{110}$$

of the Lie algebra bundle $V_G P \to X$ (93). We call ξ (110) the infinitesimal principal gauge transformations. They form a Lie $C^\infty(X)$-algebra.

Any (local) one-parameter group of principal automorphism Φ_P (108) of a principal bundle P admits the jet prolongation $J^1\Phi_P$ to a one-parameter group of G-equivariant automorphism of the jet manifold J^1P which, in turn, yields a one-parameter group of principal automorphisms Φ_C of the bundle of principal connections C (98). Its infinitesimal generator is a vector field on C, called the principal vector field on C and regarded as an infinitesimal gauge transformation of C. Thus, any infinitesimal principal gauge transformation ξ (110) on P yields a vertical vector field

$$u_\xi = (\partial_\mu \xi^r + c^r_{pq} a^p_\mu \xi^q)\partial^\mu_r \tag{111}$$

on C. Its jet prolongation onto J^1C reads

$$J^1 u_\xi = u_\xi + (\partial_{\lambda\mu}\xi^r + c^r_{pq} a^p_\mu \partial_\lambda \xi^q + c^r_{pq} a^p_{\lambda\mu}\xi^q)\partial^{\lambda\mu}_r.$$

It is readily justified that the monomorphism

$$V_G P(X) \ni \xi \to u_\xi \in \mathcal{T}(C) \tag{112}$$

obeys the equality

$$u_{[\xi,\eta]} = [u_\xi, u_\eta], \tag{113}$$

i.e., it is a monomorphism of real Lie algebras. Moreover, a glance at the expression (111) shows that the monomorphism (112) is a linear first order differential operator which sends sections of the pull-back bundle $C \underset{X}{\times} V_G P \to C$ onto sections of the vertical tangent bundle $VC \to C$. We therefore can treat vector fields (111) as infinitesimal gauge transformations depending on gauge parameters $\xi \in V_G P(X)$ [19, 20].

Let us assume that the gauge theory Lagrangian L (106) on J^1C is invariant under vertical gauge transformations. This means that vertical vector fields (111) are exact symmetries of L, i.e.,

$$\mathbf{L}_{J^1 u_\xi} L = 0. \tag{114}$$

Then the first variational formula (54) for the Lie derivative (114) takes a form

$$0 = (\partial_\mu \xi^r + c^r_{pq} a^p_\mu \xi^q)\mathcal{E}^\mu_r + d_\lambda[(\partial_\mu \xi^r + c^r_{pq} a^p_\mu \xi^q)\partial^{\lambda\mu}_r \mathcal{L})]. \tag{115}$$

It leads to the gauge invariance conditions

$$\partial^{\mu\lambda}_p\mathcal{L} + \partial^{\lambda\mu}_p\mathcal{L} = 0, \tag{116}$$

$$\mathcal{E}^\mu_r + d_\lambda\partial^{\lambda\mu}_r\mathcal{L} + c^q_{pr}a^p_\nu\partial^{\mu\nu}_q\mathcal{L} = 0, \tag{117}$$

$$c^r_{pq}(a^p_\mu\mathcal{E}^\mu_r + d_\lambda(a^p_\mu\partial^{\lambda\mu}_r\mathcal{L})) = 0. \tag{118}$$

One can regard the equalities (116) – (118) as the conditions of a Lagrangian L to be gauge invariant. They are brought into the form

$$\partial^{\mu\lambda}_p\mathcal{L} + \partial^{\lambda\mu}_p\mathcal{L} = 0. \tag{119}$$

$$\partial^\mu_q\mathcal{L} + c^r_{pq}a^p_\nu\partial^{\mu\nu}_r\mathcal{L} = 0, \tag{120}$$

$$c^r_{pq}(a^p_\mu\partial^\mu_r\mathcal{L} + a^p_{\lambda\mu}\partial^{\lambda\mu}_r\mathcal{L}) = 0. \tag{121}$$

Let us utilize the coordinates $(a^q_\mu, \mathcal{F}^r_{\lambda\mu}, \mathcal{S}^r_{\lambda\mu})$ (104) which correspond to the canonical splitting (103) of the affine jet bundle $J^1C \to C$. With respect to these coordinates, the equation (119) reads

$$\frac{\partial\mathcal{L}}{\partial\mathcal{S}^p_{\mu\lambda}} = 0. \tag{122}$$

Then the equation (120) takes a form

$$\frac{\partial\mathcal{L}}{\partial a^q_\mu} = 0. \tag{123}$$

A glance at the equalities (122) and (123) shows that a gauge invariant Lagrangian factorizes through the strength coordinates $\mathcal{F}$ (104). Then the equation (121), written as

$$c^r_{pq}\mathcal{F}^p_{\lambda\mu}\frac{\partial\mathcal{L}}{\partial\mathcal{F}^r_{\lambda\mu}} = 0,$$

shows that the gauge symmetry u_ξ of a Lagrangian L is exact. The following thus has been proved.

Theorem 20. A gauge theory Lagrangian (106) possesses the exact gauge symmetry u_ξ (111) only if it factorizes through the strength $\mathcal{F}$ (104).

A corollary of this result is the well-known Utiyama theorem [11].

Theorem 21. There is a unique gauge invariant quadratic first order Lagrangian (with the accuracy to variationally trivial ones). It is the conventional Yang–Mills Lagrangian

$$L_{\rm YM} = \frac{1}{4}a^G_{pq}g^{\lambda\mu}g^{\beta\nu}\mathcal{F}^p_{\lambda\beta}\mathcal{F}^q_{\mu\nu}\sqrt{|g|}\,\omega, \qquad g = \det(g_{\mu\nu}), \tag{124}$$

where a^G is a G-invariant bilinear form on the Lie algebra $\mathfrak{g}_r$ and g is a world metric on X.

The Euler–Lagrange operator (107) of the Yang–Mills Lagrangian $L_{\rm YM}$ (124) is

$$\mathcal{E}_{\rm YM} = \mathcal{E}^\mu_r\theta^\mu_r \wedge \omega = (\delta^n_r d_\lambda + c^n_{rp}a^p_\lambda)(a^G_{nq}g^{\mu\alpha}g^{\lambda\beta}\mathcal{F}^q_{\alpha\beta}\sqrt{|g|})\theta^r_\mu \wedge \omega. \tag{125}$$

Its kernel defines the Yang–Mills equations

$$\mathcal{E}^\mu_r = (\delta^n_r d_\lambda + c^n_{rp}a^p_\lambda)(a^G_{nq}g^{\mu\alpha}g^{\lambda\beta}\mathcal{F}^q_{\alpha\beta}\sqrt{|g|}) = 0.$$

We call a Lagrangian system $(\mathcal{S}^*_\infty[C], L_{\rm YM})$ the Yang–Mills gauge theory.

Remark 8. In classical gauge theory, there are Lagrangians, e.g., the Chern–Simons one which do not factorize through the strength of a gauge field, and whose gauge symmetry u_ξ (111) is variational, but not exact.

Since the gauge symmetry u_ξ of the Yang–Mills Lagrangian (124) is exact, the first variational formula (115) leads to the weak conservation law (55):

$$0 \approx d_\lambda(-u_\xi{}^\mu_r \partial^{\lambda\mu}_r \mathcal{L}_{\rm YM}), \tag{126}$$

of the Noether current (75):

$$\mathcal{J}^\lambda_\xi = -(\partial_\mu \xi^r + c^r_{pq} a^p_\mu \xi^q)(a^G_{rq} g^{\mu\alpha} g^{\lambda\beta} \mathcal{F}^q_{\alpha\beta} \sqrt{|g|}). \tag{127}$$

In accordance with Theorem 17, the Noether current (127) is brought into the superpotential form (76) which reads

$$\mathcal{J}^\lambda_\xi = \xi^r \mathcal{E}^\mu_r + d_\nu(\xi^r \partial^{[\nu\mu]}_r \mathcal{L}_{\rm YM}), \qquad U^{\nu\mu} = \xi^r a^G_{rq} g^{\nu\alpha} g^{\mu\beta} \mathcal{F}^q_{\alpha\beta} \sqrt{|g|}.$$

The gauge invariance conditions (116) – (118) lead to the Noether identities

$$c^p_{rq} a^q_\mu \mathcal{E}^\mu_p + d_\mu \mathcal{E}^\mu_r = 0. \tag{128}$$

which the Euler–Lagrange operator $\mathcal{E}_{\rm YM}$ (125) of the Yang–Mills Lagrangian (124) satisfies. These Noether identities are associated to the gauge symmetry u_ξ (111). One can show the following [20].

Lemma 22. The Noether identities (128) are non-trivial, complete an irreducible..

Thus, Yang–Mills gauge theory is an irreducible degenerate Lagrangian theory characterized by the complete Noether identities (128).

Following inverse second Noether Theorem 16, let us consider the differential bigraded algebra

$$P^*_\infty\{0\} = \mathcal{S}^*_\infty[\overline{VC} \underset{C}{\oplus} V_G P; C \underset{X}{\times} \overline{V_G P}] \tag{129}$$

with the local generating basis $(a^r_\mu, \overline{a}^\mu_r, c^r, \overline{c}_r)$ where c_r are odd ghosts. The gauge operator u (74) associated to the Noether identities (128) reads

$$u = (c^r_\mu + c^r_{pq} a^p_\mu c^q)\partial^\mu_r. \tag{130}$$

It is an odd gauge symmetry of the Yang–Mills Lagrangian $L_{\rm YM}$ which can be obtained from the gauge symmetry u_ξ (111) by replacement of gauge parameters ξ^r with odd ghosts c^r.

Furthermore, due to the relation (113), the gauge operator u (130) admits the nilpotent BRST extension

$$\mathbf{b} = (c^r_\mu + c^r_{pq} a^p_\mu c^q)\frac{\partial}{\partial a^r_\mu} - \frac{1}{2} c^r_{pq} c^p c^q \frac{\partial}{\partial c^r},$$

which is the well-known BRST operator in Yang–Mills gauge theory [21]. Then, by virtue of Theorem 19, the Yang–Mills Lagrangian $L_{\rm YM}$ is extended to a proper solution of the master equation

$$L_E = L_{\rm YM} + (c^r_\mu + c^r_{pq} a^p_\mu c^q)\overline{a}^\mu_r \omega - \frac{1}{2} c^r_{pq} c^p c^q \overline{c}_r \omega.$$

10 Principal graded bundles

By analogy to gauge theory on principal bundles, we develop SUSY gauge theory as theory of graded principal connections on principal graded bundles.

Graded principal bundles and connections on these bundles [25, 31, 36] can be studied similarly to principal superbundles and principal superconnections [4, 25, 31]. Since the definition of a graded Lie group and its Lie superalgebra involves the notion of a Hopf algebra, we start with the following construction [36].

Let $(Z,\mathfrak{A})$ be a graded manifold of dimension (n,m). One considers the finite dual $\mathfrak{A}(Z)^\circ$ of its structure ring $\mathfrak{A}(Z)$ which consists of elements a of the dual $\mathfrak{A}(Z)^*$ of $\mathfrak{A}(Z)$ vanishing on an ideal of $\mathfrak{A}(Z)$ of finite codimension. This is brought into a graded commutative coalgebra with the comultiplication

$$\Delta^\circ(a)(f\otimes f')=a(ff'), \qquad f,f'\in\mathfrak{A}(Z), \qquad a\in\mathfrak{A}(Z)^\circ,$$

and the counit

$$\epsilon^\circ(a)=a(\mathbf{1}_{\mathfrak{A}(Z)}).$$

In particular, $\mathfrak{A}(Z)^\circ$ includes the evaluation elements δ_z such that

$$\delta_z(f)=(\sigma(f))(z).$$

Given an evaluation element δ_z, elements $u\in\mathfrak{A}(Z)^\circ$ are said to be primitive relative to δ_z if they obey a relation

$$\Delta^\circ(u)=u\otimes\delta_z+\delta_z\otimes u.$$

These elements are derivations of $\mathfrak{A}(Z)$ at z, i.e.,

$$u(ff')=u(f)\delta_z(f')+(-1)^{[u][f]}\delta_z(f)u(f').$$

Definition 23. A graded Lie group $(G,\mathcal{G})$ is defined as a graded manifold whose body G is a real Lie group, and the structure ring $\mathcal{G}(G)$ is a graded Hopf algebra (Δ,ϵ,S) such that a body epimorphism $\sigma:\mathcal{G}(G)\to C^\infty(G)$ is a Hopf algebra morphism where a ring $C^\infty(G)$ of smooth real functions on a Lie group G possesses a real Hopf algebra structure with the co-operations:

$$\Delta(f)(g,g')=f(gg'), \qquad \epsilon(f)=f(\mathbf{1}), \qquad S(f)(g)=f(g^{-1}), \qquad f\in C^\infty(G).$$

One can show that the finite dual $\mathcal{G}(G)^\circ$ of $\mathcal{G}(G)^\circ$ is equipped with the structure of a real Hopf algebra with the multiplication law

$$a\star b=(a\otimes b)\circ\Delta, \qquad a,b\in\mathcal{G}(G)^\circ. \tag{131}$$

With respect to this multiplication, the evaluation elements δ_g, $g\in G$, provided with the product $\delta_g\star\delta_{g'}=\delta_{gg'}$ (131), constitute a group isomorphic to G. They are called the group-like elements.

It is readily observed that the set of primitive elements of $\mathcal{G}(G)^\circ$ relative to $\delta_{\mathbf{1}}$ is a real Lie superalgebra $\mathfrak{g}$ with respect to the bracket

$$[u, u'] = u \star u' - (-1)^{[u][u']} u' \star u. \tag{132}$$

It is called the Lie superalgebra of a graded Lie group $(G, \mathcal{G})$. Its even part $\mathfrak{g}_0$ is a Lie algebra of a Lie group G.

One says that a graded Lie group $(G, \mathcal{G})$ acts on a graded manifold $(Z, \mathfrak{A})$ on the right if there exists a morphism of graded manifolds

$$(\varphi, \Phi) : (Z, \mathfrak{A}) \times (G, \mathcal{G}) \to (Z, \mathfrak{A})$$

such that the corresponding ring morphism

$$\Phi : \mathfrak{A}(Z) \to \mathfrak{A}(Z) \otimes \mathcal{G}(G)$$

defines a structure of a right $\mathcal{G}(G)$-comodule on $\mathfrak{A}(Z)$, i.e.,

$$(\mathrm{Id} \otimes \Delta) \circ \Phi = (\Phi \otimes \mathrm{Id}\,) \circ \Phi, \qquad (\mathrm{Id} \otimes \epsilon) \circ \Phi = \mathrm{Id}\,.$$

For a right action (φ, Φ) and for each element $a \in \mathcal{G}(G)^\circ$, one can introduce a linear map

$$\Phi_a = (\mathrm{Id} \otimes a) \circ \Phi : \mathfrak{A}(Z) \to \mathfrak{A}(Z). \tag{133}$$

In particular, if a is a primitive element with respect to δ_e, then $\Phi_a \in \mathfrak{d}\mathfrak{A}(Z)$.

Let us consider a right action of a graded Lie group $(G, \mathcal{G})$ on itself. If $\Phi = \Delta$ and $a = \delta_g$ is a group-like element, then

$$r_g = (\mathrm{Id} \otimes \delta_g) \circ \Delta \tag{134}$$

is an even graded algebra isomorphism which corresponds to the right translation $G \to Gg$. Similarly, the left action

$$l_g = (\delta_g \otimes \mathrm{Id}\,) \circ \Delta \tag{135}$$

is defined. If $a \in \mathfrak{g}$, then l_a is a derivation of $\mathcal{G}(G)$. Given a basis $\{u_i\}$ for $\mathfrak{g}$, the derivations Φ_{u_i} constitute the global basis for $\mathfrak{d}\mathcal{G}(G)$, i.e., $\mathfrak{d}\mathcal{G}(G)$ is a free left $\mathcal{G}(G)$-module. In particular, there is a decomposition

$$\begin{aligned}
&\mathcal{G}(G) = \mathcal{G}'(G) \oplus_R \mathcal{G}''(G),\\
&\mathcal{G}'(G) = \{f \in \mathcal{G}(G) \,:\, \Phi_u(f) = 0,\ u \in \mathfrak{g}_0\},\\
&\mathcal{G}''(G) = \{f \in \mathcal{G}(G) \,:\, \Phi_u(f) = 0,\ u \in \mathfrak{g}_1\}.
\end{aligned}$$

Since $\mathcal{G}''(G) \cong C^\infty(G)$, one can show the following [1, 9, 24].

Theorem 24. A graded Lie group $(G, \mathcal{G})$ is a simple graded manifold modelled over a trivial bundle

$$G \times \wedge\mathfrak{g}_1^* \to G, \tag{136}$$

endowed with the right action r_g, $g \in G$, (134) of a group G.

Let us turn now to the notion of a principal graded bundle, but, following Section 5, we restrict our consideration to graded bundles over a smooth manifold X. A right action (φ, Φ)

of graded Lie group $(G,\mathcal{G})$ on a graded manifold $(Z,\mathfrak{A})$ is called free if, for each $z\in Z$, the morphism $\Phi_z:\mathfrak{A}(Z)\to\mathcal{G}(G)$ is such that the dual morphism $\Phi_{z*}:\mathcal{G}(G)^\circ\to\mathfrak{A}(Z)^\circ$ is injective. A right action (φ,Φ) of $(G,\mathcal{G})$ on $(Z,\mathfrak{A})$ is said to be regular if the morphism

$$(\varphi\times \mathrm{pr}_1)\circ\Delta:(Z,\mathfrak{A})\times(G,\mathcal{G})\to(Z,\mathfrak{A})\times(Z,\mathfrak{A})$$

defines a closed graded submanifold of $(Z,\mathfrak{A})\times(Z,\mathfrak{A})$.

Remark 9. Let us note that $(Z',\mathfrak{A}')$ is said to be a graded submanifold of $(Z,\mathfrak{A})$ if there exists a morphism $(Z',\mathfrak{A}')\to(Z,\mathfrak{A})$ such that the corresponding morphism $\mathfrak{A}'(Z')^\circ\to\mathfrak{A}(Z)^\circ$ is an inclusion. A graded submanifold is called closed if $\dim(Z',\mathfrak{A}')<\dim(Z,\mathfrak{A})$.

Then we come to the following variant of the well-known theorem on the quotient of a graded manifold [1, 36].

Theorem 25. A right action (φ,Φ) of $(G,\mathcal{G})$ on $(Z,\mathfrak{A})$ is regular iff the quotient $(Z/G,\mathfrak{A}/\mathcal{G})$ is a graded manifold, i.e., there exists an epimorphism of graded manifolds $(Z,\mathfrak{A})\to(Z/G,\mathfrak{A}/\mathcal{G})$ compatible with the surjection $Z\to Z/G$.

In view of this Theorem, a principal graded bundle $(P,\mathfrak{A})$ can be defined as a locally trivial submersion

$$(P,\mathfrak{A})\to(P/G,\mathfrak{A}/\mathcal{G})$$

with respect to the right regular free action of $(G,\mathcal{G})$ on $(P,\mathfrak{A})$. In an equivalent way, one can say that a principal graded bundle is a graded manifold $(P,\mathfrak{A})$ together with a free right action of a graded Lie group $(G,\mathcal{G})$ on $(P,\mathfrak{A})$ such that the quotient $(P/G,\mathfrak{A}/\mathcal{G})$ is a graded manifold and the natural surjection $(P,\mathfrak{A})\to(P/G,\mathfrak{A}/\mathcal{G})$ is a submersion. Obviously, $P\to P/G$ is a familiar principal bundle with a structure group G.

As was mentioned above, we restrict our consideration to the case of a principal graded bundle whose base a trivial graded manifold $(X=P/G,\mathfrak{A}/\mathcal{G}=C^\infty_X)$. i.e., principal graded bundles over a smooth manifold X.

Definition 26. A principal graded bundle is defined to be is a simple graded manifold $(P,\mathfrak{A}_F)$ whose body is a principal bundle $P\to X$ with a structure group G, and which is modelled over a composite bundle

$$F=P\underset{X}{\times}W\to P\to X \tag{137}$$

where $W\to X$ is P-associated bundle with a typical fibre $\mathfrak{g}_1$ provided with the left action (135) of a group G. A principal graded bundle $(P,\mathfrak{A}_F)$ is subject to the right fibrewise action (133) of a group G.

Remark 10. Being kept as a simple graded manifold, a graded bundle $(P,\mathfrak{A}_F)$ fails to admit an action of a graded Lie group $(G,\mathcal{G})$. Therefore, it is not a principal graded bundle in a strict sense.

11 Yang–Mills supergauge theory

Let $\mathfrak{g}$ be a finite-dimensional real Lie superalgebra with the basis $\{\varepsilon_r\}$, $r = 1, \ldots, m$, and real structure constants c^r_{ij} such that

$$c^r_{ij} = -(-1)^{[i][j]}c^r_{ji}, \qquad [r] = [i] + [j],$$
$$(-1)^{[i][b]}c^r_{ij}c^j_{ab} + (-1)^{[a][i]}c^r_{aj}c^j_{bi} + (-1)^{[b][a]}c^r_{bj}c^j_{ia} = 0,$$

where $[r]$ denotes the Grassmann parity of ε_r. Its even part $\mathfrak{g}_0$ is a Lie algebra of some Lie group G. Let the adjoint representation of $\mathfrak{g}_0$ in $\mathfrak{g}$ is extended to the corresponding action of G on $\mathfrak{g}$.

Let P be a principal bundle with a structure group G. Given Yang–Mills gauge theory of principal connections on P (see Section 9), we aim to extend it to a Yang–Mills supergauge theory associated to a Lie superalgebra $\mathfrak{g}$.

Let us consider a simple graded manifold $(G, \mathcal{G})$ modelled over the trivial vector bundle (136). In accordance with Theorem 24, this is a graded Lie group whose Lie superalgebra is $\mathfrak{g}$. Following Definition 26, let us define a principal graded bundle $(P, \mathfrak{A}_F)$ modelled over the composite bundle (137).

Let J^1F be a jet manifold of the fibre bundle $F \to X$ (137). It is a composite bundle

$$J^1F \to J^1P \to X, \tag{138}$$

where $J^1F \to J^1P$ is a vector bundle. Then a simple graded manifold (J^1P, A_{J^1f}) modelled over $J^1F \to J^1P$ is a graded jet manifold of a principal graded bundle $(P, \mathfrak{A}_F)$ in accordance with Definition 7. The composite bundle (138) is provided with the jet prolongation of the right action (133) of G onto $(P, \mathfrak{A}_F)$. By analogy with the bundle of principal connections C (98), let us consider the quotient

$$\mathcal{C} = J^1F/G \to C \to X. \tag{139}$$

Since $\mathcal{C} \to C$ is a vector bundle, we have a simple graded manifold $(C, \mathfrak{A}_\mathcal{C})$ modelled over a vector bundle $\mathcal{C} \to C$. It is a graded bundle over a smooth manifold X.

We develop supergauge theory as a Lagrangian theory on a graded bundle $(C, \mathfrak{A}_\mathcal{C})$ (see Section 7). Given a basis $\{\varepsilon_r\}$ for $\mathfrak{g}$, we have a local basis (x^λ, a^r_λ) of Grassmann parity $[a^r_\lambda] = [r]$ for a graded bundle $(C, \mathfrak{A}_\mathcal{C})$. Similarly to the splitting (104), jets of the elements a^r_λ admit the decomposition

$$a^r_{\lambda\mu} = \frac{1}{2}(\mathcal{F}^r_{\lambda\mu} + \mathcal{S}^r_{\lambda\mu}) = \tag{140}$$
$$\frac{1}{2}(a^r_{\lambda\mu} - a^r_{\mu\lambda} + c^r_{ij}a^i_\lambda a^j_\mu) + \frac{1}{2}(a^r_{\lambda\mu} + a^r_{\mu\lambda} - c^r_{ij}a^i_\lambda a^j_\mu).$$

Let us consider the differential bigraded algebra $\mathcal{S}^*_\infty[\mathcal{C}; C]$ (41). Given the universal enveloping algebra $\overline{\mathfrak{g}}$ of $\mathfrak{g}$, we assume that there exists an even quadratic Casimir element $h^{ij}\varepsilon_i\varepsilon_j$ of $\overline{\mathfrak{g}}$ such that the matrix h^{ij} is non-degenerate. Let g be metric on X. Then a graded Yang–Mills Lagrangian is defined as

$$L_{\rm YM} = \frac{1}{4}h_{ij}g^{\lambda\mu}g^{\beta\nu}\mathcal{F}^i_{\lambda\beta}\mathcal{F}^j_{\mu\nu}\omega.$$

Its variational derivatives $\mathcal{E}^\lambda_r$ obey the irreducible complete Noether identities

$$c^r_{ji}a^i_\lambda\mathcal{E}^\lambda_r + d_\lambda\mathcal{E}^\lambda_j = 0.$$

Therefore, let us enlarge the differential bigraded algebra $\mathcal{S}^*_\infty[\mathcal{C};C]$ to the differential bigraded algebra

$$P^*_\infty\{0\} = \mathcal{S}^*_\infty[\mathcal{C}\underset{X}{\oplus}\overline{\mathcal{C}},\underset{X}{\oplus}VF/G\underset{X}{\oplus}\overline{VF/G};P],$$

where $\overline{E}$ denotes the density dual (58) of a fibre bundle $E\to X$. Its local basis $(x^\lambda, a^r_\lambda, \overline{a}^\lambda_r, c^r, \overline{c}_r)$ contains gauge fields a^r_λ, their antifields $\overline{a}^\lambda_r$ of Grassmann parity

$$[\overline{a}^\lambda_r] = ([r]+1)\mathrm{mod}\,2,$$

the ghosts c^r of Grassmann parity

$$[c^r] = ([r]+1)\mathrm{mod}\,2,$$

and the Noether antifields $\overline{c}_r$ of Grassmann parity $[\overline{c}_r]=[r]$. Then the gauge operator (74) reads

$$u = (c^r_\lambda - c^r_{ji}c^j a^i_\lambda)\frac{\partial}{\partial a^r_\lambda} \tag{141}$$

(cf. (130)). It admits the nilpotent BRST extension

$$\mathbf{b} = (c^r_\lambda - c^r_{ji}c^j a^i_\lambda)\frac{\partial}{\partial a^r_\lambda} - \frac{1}{2}(-1)^{[i]}c^r_{ij}c^i c^j\frac{\partial}{\partial c^r}.$$

The corresponding proper solution (83) of the master equation takes a form

$$L_E = L_{\mathrm{YM}} + (c^r_\lambda - c^r_{ji}c^j a^i_\lambda)\overline{a}^\lambda_r\omega - \frac{1}{2}(-1)^{[i]}c^r_{ij}c^i c^j\overline{c}_r\omega.$$

References

[1] A. Almorox, Supergauge theories in graded manifolds, In: *Differential Geometric Methods in Mathematical Physics*, Lect. Notes in Math. **1251** (Springer, Berlin, 1987), p.114.

[2] I. Anderson and T. Duchamp, On the existence of global variational principles, *Amer. J. Math.* **102** (1980) 781.

[3] G. Barnich, F. Brandt, M. Henneaux, Local BRST cohomology in gauge theories, *Phys. Rep.* **338** (2000) 439.

[4] C. Bartocci, U. Bruzzo and D. Hernández Ruipérez, *The Geometry of Supermanifolds* (Kluwer, Dordrecht, 1991).

[5] D. Bashkirov, G. Giachetta, L. Mangiarotti and G. Sardanashvily, Noether's second theorem for BRST symmetries, *J. Math. Phys.* **46** (2005) 053517; *arXiv*: math-ph/ 0412034.

[6] D. Bashkirov, G. Giachetta, L. Mangiarotti and G. Sardanashvily, The antifield Koszul–Tate complex of reducible Noether identities, *J. Math. Phys.* **46** (2005) 103513; *arXiv*: math-ph/ 0506034.

[7] D. Bashkirov, G. Giachetta, L. Mangiarotti and G. Sardanashvily, The KT-BRST complex of degenerate Lagrangian systems, *Lett. Math. Phys.* **83** (2008) 237; *arXiv*: math-ph/ 0702097.

[8] M. Batchelor, The structure of supermanifolds, *Trans. Amer. Math. Soc.* **253** (1979) 329.

[9] C. Boyer and O. Sánchez Valenzuela, Lie supergroup action on supermanifolds, *Trans. Amer. Math. Soc.* **323** (1991) 151.

[10] F. Brandt, Jet coordinates for local BRST cohomology, *Lett. Math. Phys.* **55** (2001) 149.

[11] U. Bruzzo, The global Utiyama theorem in Einstein–Cartan theory, *J. Math. Phys.* **28**, (1987) 2074.

[12] J. Cariñena, H. Figueroa, Singular Lagrangian in supermechanics, *Diff. J. Geom. Appl.* **18** (2003) 33.

[13] R. Cianci, M. Francaviglia, I. Volovich, Variational calculus and Poincaré–Cartan formalism in supermanifolds, *J. Phys. A.* **28** (1995) 723.

[14] D. Franco, C. Polito, Supersymmetric field-theoretic models on a supermanifold, *J. Math. Phys.* **45** (2004) 1447.

[15] D. Fuks, *Cohomology of Infinite-Dimensional Lie Algebras* (Consultants Bureau, New York, 1986).

[16] D. Hernández Ruipérez and J. Muñoz Masqué, Global variational calculus on graded manifolds, *J. Math. Pures Appl.* **63** (1984) 283.

[17] G. Giachetta, L. Mangiarotti, and G. Sardanashvily, *Geometric and Algebraic Topological Methods in Quantum Mechanics* (World Scientific, Singapore, 2005).

[18] G. Giachetta, L. Mangiarotti and G. Sardanashvily, Lagrangian supersymmetries depending on derivatives. Global analysis and cohomology, *Commun. Math. Phys.*, **259** (2005) 103.

[19] G. Giachetta, L. Mangiarotti and G. Sardanashvily, On the notion of gauge symmetries of generic Lagrangian field theory, *J. Math. Phys* **50**(2009) 012903; *arXiv*: arXiv: 0807.3003.

[20] G. Giachetta, L. Mangiarotti, and G. Sardanashvily, *Advanced Classical Field Theory* (World Scientific, Singapore, 2009).

[21] J. Gomis, J. París and S. Samuel, S. (1995). Antibracket, antifields and gauge theory quantization, *Phys. Rep.* **295** (1995) 1.

[22] A. Jadczyk and K. Pilch, (1981). Superspaces and Supersymmetries, *Commun. Math. Phys.* **78** (1981) 391.

[23] S. Kobayashi and K. Nomizu, *Foundations of Differential Geometry* (John Wiley, New York - Singapore, 1963).

[24] B. Kostant, Graded manifolds, graded Lie theory, and prequantization, In: *Differential Geometric Methods in Mathematical Physics*, Lect. Notes in Math. **570** (Springer, Berlin, 1977) p. 177.

[25] L. Mangiarotti and G. Sardanashvily, *Connections in Classical and Quantum Field Theory* (World Scientific, Singapore, 2000).

[26] J. Monterde, J. Munos Masque, Variational problems on graded manifolds, *Contemp. Math.* **132** (1992) 551.

[27] J. Monterde, J. Munos Masque, J. Vallejo, The Poincare–Cartan form in superfield theory, *Int. J. Geom. Methods Mod. Phys.* **3** (2006) 775.

[28] A. Rennie, Smoothness and locality for nonunital spectral triples, *K-Theory* **28** (2003) 127.

[29] G. Sardanashvily, Graded infinite order jet manifolds, *Int. J. Geom. Methods Mod. Phys.* **4**, (2007) 1335; *arXiv*: arXiv: 0708.2434.

[30] G. Sardanashvily, Gauge conservation laws in a general setting: Superpotential, *Int. J. Geom. Methods Mod. Phys.* **6** (2009) 1047; *arXiv*: arXiv: 0906.1732.

[31] G. Sardanashvily, Lectures on supergeometry, *arXiv*: 0910.0092.

[32] G. Sardanashvily, *Lectures on Differential Geometry of Modules and Rings. Application to Quantum Theory* (Lambert Academic Publishing, Saarbrucken, 2012); *arXiv*: 0910.1515.

[33] G. Sardanashvily, *Advanced Differential Geometry for Theoreticians. Fiber bundles, jet manifolds and Lagrangian theory* (Lambert Academic Publishing, Saarbrucken, 2013); *arXiv*: 0908.1886.

[34] G. Sardanashvily, Graded Lagrangian formalism, *Int. J. Geom. Methods. Mod. Phys.* **10** (2013) 1350016; *arXiv*: 1206.2508.

[35] G. Sardanashvily, Remark on the Serre – Swan theorem fort graded manifolds, *arXiv*: 1304.1371.

[36] T. Stavracou, Theory of connections on graded principal bundles, *Rev. Math. Phys.* **10** (1998) 47.

[37] F. Takens, A global version of the inverse problem of the calculus of variations, *J. Diff. Geom.* **14** (1979) 543.

[38] B. Tennison, *Sheaf Theory* (Cambridge Univ. Press, Cambridge, 1975).